

Citation for published version:

Margos, G, Vollmer, SA, Ogden, NH & Fish, D 2011, 'Population genetics, taxonomy, phylogeny and evolution of *Borrelia burgdorferi sensu lato*', *Infection, Genetics and Evolution*, vol. 11, no. 7, pp. 1545-1563.
<https://doi.org/10.1016/j.meegid.2011.07.022>

DOI:

[10.1016/j.meegid.2011.07.022](https://doi.org/10.1016/j.meegid.2011.07.022)

Publication date:

2011

Document Version

Peer reviewed version

[Link to publication](#)

NOTICE: this is the author's version of a work that was accepted for publication in *Infection, Genetics and Evolution*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Infection, Genetics and Evolution*, vol 11, issue 2, 2011, DOI 10.1016/j.meegid.2011.07.022

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Population genetics, taxonomy, phylogeny and evolution of *Borrelia*
burgdorferi sensu lato

Gabriele Margos^{1*}, Stephanie A. Vollmer¹, Nicholas H. Ogden², Durland Fish³

¹Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath
BA2 7AY, UK

²Zoonoses Division, Centre for Food-borne, Environmental and Zoonotic Infectious
Diseases, Public Health Agency of Canada, Ottawa, Canada

³Department of Epidemiology and Public Health, Yale School of Medicine, Yale
University, New Haven, CT 06520

*corresponding author:

Department of Biology and Biochemistry

3 South

University of Bath

Claverton Down

Bath BA2 7AY

U.K.

Telefon: +44-1225-385116

Fax: +44-1225-38

e-mail gm250@bath.ac.uk

Abstract

To understand the population structure and dynamics of bacterial microorganisms, typing systems that accurately reflect the phylogenetic and evolutionary relationship of the agents are required. Over the past 15 years multilocus sequence typing schemes have replaced single locus approaches, giving novel insights into phylogenetic and evolutionary relationships of many bacterial species and facilitating taxonomy. Since 2004, several schemes using multiple loci have been developed to better understand the taxonomy, phylogeny and evolution of Lyme borreliosis spirochetes and in this paper we have reviewed and summarized the progress that has been made for this important group of vector-borne zoonotic bacteria.

KEYWORDS: *BORRELIA BURGDORFERI*, EVOLUTION, PHYLOGENY, MOLECULAR ECOLOGY, IXODES, TICKS, MULTILOCUS SEQUENCE TYPING, MLST

ABBREVIATIONS:

IGS – INTERGENIC SPACER

LB – LYME BORRLIOSIS

MLST/MLSA – MULTILOCUS SEQUENCE TYPING/MULTILOCUS SEQUENCE ANALYSIS

MW – MIDWEST

NE – NORTHEAST

OSP – OUTER SURFACE PROTEIN

SLV/DLV/TLV – SINGLE LOCUS VARIANT/DOUBLE LOCUS VARIANT/TRIPLE LOCUS VARIANT

SNP – SINGLE NUCLEOTIDE POLYMORPHISM

ST – SEQUENCE TYPE

1. INTRODUCTION

Tick-borne diseases are of increasing public health concern because of range expansions of both vectors and pathogens (Daniel et al., 2003; Falco et al., 1995; Ogden et al., 2008b). To understand these processes and to predict future trajectories, detailed data on the contemporary population structure and on the evolutionary and demographic histories that have shaped the populations are essential. Population structure, evolutionary and demographic processes of microbial pathogens may best be inferred using genetic data with neutral variation. Such data together with information on host associations are critical to understand the dynamics of tick-borne disease agents and to form hypothesis concerning past and future spread.

Lyme borreliosis (LB) is the most prevalent vector-borne disease in the Holarctic region (Dennis and Hayes, 2002). Due to the pattern and breadth of the ecological niches occupied by its members, the LB group of spirochetes constitutes an ideal system to investigate the contributions of host and vectors in pathogen demographic processes. In addition, major advances in sequencing technologies and the development of sophisticated typing tools for bacterial pathogens have greatly enhanced the potential to infer robust phylogenies and to deduce more accurately the evolutionary relationships of micro-organisms. In particular, targeted gene amplification and sequence analysis of several housekeeping genes, termed multilocus sequence typing or multilocus sequence analysis (MLST/MLSA) and, more recently, genome-wide detection of single nucleotide polymorphisms (SNPs) have made major contributions to advancing knowledge in bacterial population genetics, phylogenetics and molecular taxonomy (Aanensen and Spratt, 2005; Bishop et al., 2009; Hall, 2007; Harris et al., 2010; Holt et al., 2008; Maiden, 2006). In this review we are focussing

on progress that has been made in recent years using molecular methods including MLST and MLSA to study population genetics, molecular taxonomy, phylogenetics and the evolution of the LB group of spirochetes (also referred to as *Borrelia burgdorferi* sensu lato (s.l.) species complex). Although we acknowledge that not all species belonging to this species complex cause LB, we prefer to use the term ‘LB group of spirochetes’ (instead of *B. burgdorferi* s.l.) to refer to the whole group as this simplifies distinguishing *B. burgdorferi* s.l. from *B. burgdorferi* sensu stricto (the species to which we will refer hereafter as *B. burgdorferi*).

The species complex currently consists of 18 proposed and confirmed species (Margos et al., 2010; Rudenko et al., 2009a; Rudenko et al., 2009b) (Table 1), several of which can cause LB in humans (or Lyme disease). LB species vary in their geographic distribution, host specificity and ability to cause disease in humans. Clinically the different pathogenic *Borrelia* spp. are of interest as they have been associated with different disease symptoms which may be observed in the late stages of the condition. For example, *B. afzelii* is most frequently linked with skin manifestations, *B. garinii* and *B. bavariensis* with neuroborreliosis, and *B. burgdorferi* with arthritic symptoms (Canica et al., 1993; Ornstein et al., 2001; Randolph, 2008; Rijpkema et al., 1997; Stanek and Strle, 2009; Steere et al., 1986; van Dam, 2002). Some species, such as *B. lusitaniae*, have only occasionally been associated with human disease while for others, such as *B. valaisiana*, the status is uncertain because they have high regional prevalence in Europe but have rarely been isolated from humans (Collares-Pereira et al., 2004; Diza et al., 2004). It has been suggested that not all strains/genotypes within a species cause disseminated disease (Baranton et al., 2001; Seinost et al., 1999; Wilske et al., 1996; Wilske et al., 1993; Wormser et al.,

2008) and it is, therefore, of epidemiological and clinical relevance to identify the geographic range of LB species and the spatial distributions of their genotypes.

2. Ecology of LB group of spirochetes

Due to the obligate parasitic lifestyle of LB spirochetes, their biology is intimately linked to that of their invertebrate and vertebrate hosts which also broadly defines their ecological niches (Kurtenbach et al., 2002b). The ecological niche diversity of different species varies in the degree of specialization (from generalist to specialised strategies) in terms of host and vector adaptation and this influences the geographic distribution at species and population levels. There are several excellent recent reviews regarding the ecology of LB spirochetes, describing in detail host and vector interactions (Gern, 2008; Gern and Humair, 2002; Kurtenbach et al., 2006; Masuzawa, 2004; Piesman and Gern, 2004; Tsao, 2009). Here we will only briefly describe the general ecology of LB spirochetes.

The life cycle of the LB group of spirochetes is a dynamic interplay between bacteria, reservoir hosts and vectors which is confounded by landscape and climatic factors impacting host and vector ecology (Figure 1, (Kurtenbach et al., 2006)). All known vectors of LB spirochetes belong to the genus *Ixodes* and these ticks are three host ticks, i.e. they have three feeding stages (larvae, nymphs and adult females) each utilizing a different host, although not necessarily a different host species. Except in the case of nidicolous (nest-living) tick vectors, adult female ticks prefer large animals, such as deer, as hosts which are considered not susceptible to *Borrelia* infection (Telford et al., 1988). The preference of both immature stages for small to medium sized vertebrates (mammals, birds or lizards) is essential for maintaining the bacteria in its natural transmission cycles. The bacteria are taken up during a

1 bloodmeal from an infected and infectious host, are maintained transstadially during
2 the moulting process and are then transmitted to other hosts during the subsequent
3 bloodmeal during the next life stage (Gern and Humair, 2002). Other means of
4 transmission are co-feeding transmission (between neighbouring ticks feeding on a
5 susceptible or non-susceptible host) (Ogden et al., 1997) and transovarial transmission
6 ((Gern and Humair, 2002) and references therein), although the latter may depend on
7 the tick species as it has not been experimentally demonstrated for *I. scapularis* and *I.*
8 *persulcatus* (Nefedova et al., 2004; Patrican, 1997). However, relapsing-fever like
9 spirochetes (e.g. *B. miyamotoi*) are transmitted transovarially in *Ixodes* ticks and occur
10 sympatrically with LB group spirochetes, which may explain some or perhaps all
11 observations of transovarial transmission (Piesman, 2002; Scoles et al., 2001).

12 The main vectors transmitting LB spirochetes to humans are members of the
13 *Ixodes persulcatus* species complex and are generalist feeders (i.e. they have a wide
14 host range) that follow an ambush strategy for host seeking and are widely distributed
15 in the environment (Balashov, 1972; Loye and Lane, 1988; Xu et al., 2003). These are
16 *I. ricinus* in Europe, *I. persulcatus* in Eastern Europe and Asia, *I. scapularis* and *I.*
17 *pacificus* in North America. Other experimentally confirmed vector-competent *Ixodes*
18 species are nidicolous to varying degrees, i.e. they reside in the burrows of their hosts,
19 and, having a more restricted host preference, rarely bite humans (see review by
20 (Eisen and Lane, 2002)). This raises the question as to whether the LB spirochetes
21 transmitted by nidicolous vectors are non-pathogenic for humans, or whether they are
22 pathogenic but rarely cause disease because the ticks that transmit them rarely
23 encounter humans.

24 More than 100 vertebrate species have been identified that can act as reservoir
25 hosts for LB spirochetes including rodents (wood mice, wood rats, voles, dormice,

1 squirrels, chipmunks, rats), insectivores (shrews, hedgehogs), racoons and several
2 bird species (Masuzawa, 2004; Piesman and Gern, 2004). For other species such as
3 foxes or badgers only limited information is available and it is uncertain whether they
4 constitute reservoir hosts (Gern and Sell, 2009; Matuschka et al., 2000; Miyamoto and
5 Masuzawa, 2002), although domestic dogs have been reported to be reservoir
6 competent (Mather et al., 1994). Not all vertebrate hosts are permissive or equally
7 efficient as reservoir hosts for all *Borrelia* species. The basic reproduction number R_0
8 serves as a measure of fitness of different LB group species in different host-tick
9 communities (reviewed by (Randolph, 1998) and (Tsao, 2009)). For some LB group
10 species only certain host species are able to support completion of the entire
11 transmission cycle (from a vector tick through the host to the next vector tick)
12 (Kurtenbach et al., 2002a). In Europe, these ‘host associations’ have been well studied
13 and they are an important component of the ecology of LB spirochete species. Most
14 of the LB group species in Europe are transmitted by the generalist tick, *I. ricinus*,
15 which feeds on birds as well as on rodents or other medium sized mammals, rendering
16 the tick a ‘mixing vessel’ for different strains and species. Therefore, the host
17 associations described in Europe are not driven by adaptation to an endophilic tick
18 with a narrow host preference but are truly host driven (Humair and Gern, 2000;
19 Kurtenbach et al., 1998b).. Several lines of evidence support the notion of host
20 association: 1) Experimental evidence has shown that these host associations match
21 the ability of LB group species to deflect complement mediated lysis of the
22 corresponding reservoir hosts (Kurtenbach et al., 2002b; Kurtenbach et al., 1998a;
23 Lane and Quistad, 1998; Ullmann et al., 2003). 2) *B. afzelii* and *B. bavariensis* have
24 been shown to be transmitted through rodents while *B. garinii* and *B. valaisiana* are
25 transmitted through avian reservoir hosts (Dubska et al., 2009; Hanincova et al.,

2003a; Hanincova et al., 2003b; Hu et al., 1997; Hu et al., 2001; Humair et al., 1998; Humair et al., 1999; Kurtenbach et al., 1998a; Taragel'ova et al., 2008). This does not mean that *B. garinii* infections cannot be found in mice, as bird adapted outer surface protein A (OspA) serotype 6 strains have been found in internal organs of *Apodemus* mice, but these strains are not transmitted to vector competent ticks feeding on such infected mice and, therefore, represent dead-end hosts (Kurtenbach et al., 1998a; Kurtenbach et al., 2002a) . Mechanisms permitting the transmission to hosts of such host complement incompatible LB spirochete species have been suggested (Kurtenbach et al., 2002a). 3) Recent evidence supports the view that host associations substantially shape *Borrelia* populations by impacting their migration patterns and geographical distributions (Kurtenbach et al., 2006; Vollmer et al., 2011).

The compatibility of spirochetes with tick vectors has not been studied in so much detail. While many *Ixodes* tick species are able to transmit several species of *Borrelia* (Table 1), it seems that certain *Borrelia*-vector associations are not compatible or less efficient (e.g. (Dolan et al., 1998; Masuzawa et al., 2005)). Thus, vector competence - or the lack thereof - has implications on the geographic distribution of these species (see geographic distribution). .

Consequently, *Borrelia* populations are shaped by the dynamics and demographic processes of host and vector populations, host and vector immune responses and extrinsic abiotic factors (e.g. temperature, climate, landscape connecivity) affecting host and vector populations and contact between them which together determine R_0 for each species and strain of the bacterium (Figure 1). Diversity in *Borrelia* populations arises by mutation, recombination, drift and natural selection. It has been suggested that mutation rates are low as *Borrelia* are very slow growing bacteria (Hoen et al., 2009). Genetic drift may also predominate when

effective population sizes, N_e , are small (as has been suggested for *B. burgdorferi* (Qiu et al., 2002)), this may weaken natural selection and introduce stochastic effects into allele frequencies of populations (Page and Holmes, 1998). The signature of all these processes can be inferred from genetic information obtained from present day samples but as different processes can lead to similar effects, caution needs to be exercised when interpreting data (Frank, 2002).

3. Typing tools for the LB group of spirochetes

When *B. burgdorferi* was discovered and described, it was assumed to be a single species (Burgdorfer et al., 1982; Johnson et al., 1984). The use of genome fingerprinting and other methods soon showed that the bacteria were highly diverse and in fact represented a species complex (Liveris et al., 1995; Marconi and Garon, 1992; Mathiesen et al., 1997; Postic et al., 1994; Wilske et al., 1991). Phenotypic typing tools were developed to reveal intraspecific diversity which included serotyping or multilocus enzyme electrophoresis (MLEE) (Boerlin et al., 1992; Wilske et al., 1991; Wilske et al., 1996; Wilske et al., 1995; Wilske et al., 1993). This topic is reviewed excellently by Wang and co-authors (Wang et al., 1999b) and here we concentrate on single and multilocus sequence analyses.

Sequences of single gene loci have been popular for ecological, population, epidemiological and evolutionary studies of the LB group of spirochetes. Many different genes and loci have been targeted in studies depending on the level of variation and the discriminatory power required and which species were being investigated. These included intergenic spacer (IGS) regions, the *rrs* (16S rRNA) locus, the plasmid located genes encoding the outer surface proteins A and C (*ospA*, *ospC*), decorin-binding protein A (*dbpA*), the chromosomally located housekeeping

genes recombinase A (*recA*), *groEL*, *hbb* or flagellin B (*flaB*) (Casati et al., 2004; Dykhuizen and Baranton, 2001; Fukunaga et al., 1996c; Liveris et al., 1995; Marconi et al., 1995; Michel et al., 2004; Park et al., 2004; Postic et al., 1994; Schulte-Spechtel et al., 2006; Valsangiacomo et al., 1997; Will et al., 1995; Wilske et al., 1996).

3.1 Interspecies Studies

For species definition and evolutionary studies, conserved loci or intergenic spacer have been employed. *flaB* has been popular for evolutionary studies and species identification because the *flaB* gene is present in relapsing fever spirochetes, which can be used as an outgroup to root phylogenetic trees. This conserved locus was used to create an early and reasonably complete evolutionary tree of the LB group of spirochetes (Fukunaga et al., 1996c). Some groups now use this and other conserved loci (e.g. 23S, *hbb*) to screen field-collected questing ticks (by real-time or conventional PCR) and to establish infection prevalences with LB species, as well as with relapsing-fever like spirochaete species such as *B. miyamotoi* which infects hard bodied ixodid ticks worldwide (Barbour et al., 2010; Fukunaga et al., 1995; Herrmann and Gern, 2010; Ogden et al., 2011; Portnoi et al., 2006).

The region encoding the ribosomal RNAs (rRNA) has been popular in studies of LB spirochaetes where different regions have been used for various purposes and species. The 16S (*rrs*) subunit, has been used in evolutionary and speciation studies (e.g. (Fukunaga et al., 1996a; Le Fleche et al., 1997).

Approximately 2 kb downstream of a single copy of the 16S rRNA small subunit are tandemly repeated copies of the 23S-5S (*rrl-rrf*) large subunits (Schwartz et al., 1992). The IGS between the 5S and 23S (*rrf-rrl*) of the repeated pairs is approximately 200-250 bp and this organization of rRNA genes appears to be unique

to the LB group spirochetes (Gazumyan et al., 1994; Schwartz et al., 1992). The 5S-23S spacer is possibly the most common sequence-based method for LB group species identification in Europe and approaches have recently been developed using quantitative PCR to screen questing ticks (Postic et al., 1994; Postic et al., 1998; Strube et al., 2010). Diversity at this locus has also been investigated using reverse line blot, a key method in epidemiological studies of LB species due to it being a rapid and reliable method for detecting and typing mixed infections of different *Borrelia* species in field-collected tick or host samples. It uses PCR products of the 5S-23S IGS region for hybridization to membrane bound oligonucleotides that are specific for different LB group species. This method was first used to identify the prevalence of different LB group species in ticks in The Netherlands (Rijpkema et al., 1997). Reverse line blot was better suited than some other methods for characterising mixed species infections and partly for this reason it was a key method in identifying the patterns of host specialization (Hanincova et al., 2003b; Kurtenbach et al., 2001; Kurtenbach et al., 1998a). A problem was that this method was unable to distinguish ecotypes of *B. garinii* (bird or rodent associated which are now considered different species, (Margos et al., 2009)) which may have confused some conclusions of host associations.

3.2 Intraspecies Genotyping

For intraspecies studies loci that provide good level of polymorphism have been widely used, such as the 16S-23S (*rrs-rrl*) IGS or outer surface protein (*osp*) encoding loci for *B. burgdorferi* in North America (Bunikis et al., 2004; Girard et al., 2009; Hamer et al., 2010; Hanincova et al., 2008a; Liveris et al., 1995; Marconi et al., 1995;

Ogden et al., 2008b; Postic et al., 1994). However, these are not necessarily useful for species identification or for intraspecies studies of other LB group species.

Outer surface proteins are variable and have, for this reason, often been used for population studies. *ospA*, located on a 49- to 70- kb linear plasmid, called lp54 in *B. burgdorferi* (Barbour and Garon, 1988), revealed differences in the levels of homogeneity of LB species. It was observed that there is great variation in *ospA* in *B. garinii* while there is much homogeneity in some other species such as *B. burgdorferi* or *B. afzelii* which is consistent with serotyping studies (Wilske et al., 1996). This locus has also been used to reveal rare horizontal gene transfer between species (Rosa et al., 1992; Wang et al., 2000).

ospC is located on a 26-kb circular plasmid (Sadziene et al., 1993) and has been described as the locus with the highest degree of variation (Jauris-Heipke et al., 1995; Qiu et al., 2004; Theisen et al., 1993). This locus is rarely used for species determination because, while there may be species specific motifs (Fukunaga and Hamase, 1995; Jauris-Heipke et al., 1995), recombination and plasmid exchange means that strains of the same species do not always cluster monophyletically in phylogenies (Kurtenbach et al., 2002a; Lin et al., 2002; Margos et al., 2009) (Figure 2B). However, due to the high level of variation, *ospC* has been frequently used in population studies within species, most notably within *B. burgdorferi* (Barbour and Travinsky, 2010; Hanincova et al., 2008a; Marti Ras et al., 1997; Qiu et al., 2002), and the study of *ospC* may be useful in identifying ecological traits such as host-species associations (Ogden et al., 2011) as its expression is important for tick-to-host transmission (Piesman and Schwan, 2010).

Population genetics studies on *B. burgdorferi* in the Northeastern (NE) USA have suggested that *ospA* is in linkage disequilibrium with *ospC*, a gene on a different

1 plasmid, and the 16S-23S IGS (Qiu et al., 1997) while more recent studies have
2 shown that this may be related to the spatial scale of sampling as geographic variation
3 in linkage pattern were found (Hellgren et al., 2011; Travinsky et al., 2010). In
4 addition, horizontal transfer has been demonstrated for many plasmid-encoded loci,
5 whole plasmids and also for genes on the main chromosome although it needs to be
6 emphasized that these are likely to be rare events (Barbour and Travinsky, 2010; Qiu
7 et al., 2004; Vitorino et al., 2008; Wang et al., 1999a) unpublished). While Qiu and
8 co-authors (Qiu et al., 2004), using almost exclusively plasmid-located loci, found a
9 higher rate of recombination than mutation (ratio 3:1), studies using chromosomally
10 located housekeeping genes found higher mutation than recombination rates with an
11 r/m of 1:100 to 1:25, strongly suggesting that the linear chromosome is well suited for
12 studies investigating evolutionary and population relationships of LB spirochetes
13 ((Vitorino et al., 2008), Vollmer et al. unpublished).

14 Different loci tended to be preferred in North America or Asia compared to
15 Europe. Many studies conducted in the USA, where *B. burgdorferi* is the only species
16 causing human disease, have focused on *ospC*, the 16S-23S IGS or a combination of
17 these and additional loci (Brisson and Dykhuizen, 2004; Brisson et al., 2010; Bunikis
18 et al., 2004; Girard et al., 2009; Hanincova et al., 2008a; Liveris et al., 1995; Marti
19 Ras et al., 1997; Ogden et al., 2008b; Qiu et al., 2002; Wang et al., 1999; Wormser et
20 al., 1999). In Europe several *Borrelia* species are prevalent and all four major disease-
21 causing species (i.e. *B. afzelii*, *B. garinii*, *B. burgdorferi* and *B. bavariensis*) are
22 endemic in populations of *I. persulcatus*-group ticks (Gern, 2008). For this reason
23 species definition has been the key for epidemiological and ecological studies, and
24 thus, in Europe *ospA* and the 5S-23S IGS region have been most commonly used
25 (summarized by (Rauter and Hartung, 2005)). In Asia, species identification was often

the major aim of studies and a variety of loci have been used including loci favoured in Europe as well as more conserved loci, such as *flaB* and 16S rRNA (Masuzawa, 2004). This is most likely because fewer population genetic studies have been completed and species prevalence is of primary importance over such a large area considering the broad spectrum of species found across the continent.

3.3 Typing schemes using multiple loci

Since 2004 several multilocus schemes have been developed to investigate the phylogenetic relationship of the LB spirochetes. The greater amount of genetic information obtained from several loci permits determination of more subtle differences in and between species.

MLST schemes were originally designed to utilise regions of housekeeping genes that evolved at a moderate speed to capture the intermediate relationship within bacterial species (Figure 3) (Maiden, 2006; Maiden et al., 1998). While this means that the number of polymorphic sites per gene region is usually low, by combining multiple loci the discriminatory power is high. Traditionally, internal fragments of housekeeping genes, approximately 450-500 bp long, were selected and kept in-frame. The genes were chosen throughout the genome to avoid any local bias that may occur in the bacterial genome. Another criterion was that the chosen housekeeping genes should also be flanked by genes known to have similar functions as there may be linkage between adjacent genes. If genes next to the selected housekeeping gene are under strong selection pressures, this may influence the neighbouring genes. Finally, genes should have a similar level of genetic diversity so that each gene provides a similar contribution to phylogenetic analyses and no single

gene dominates a tree generated by use of the concatenated sequences of the selected housekeeping genes (Urwin and Maiden, 2003).

One central problem when attempting to understand relationships among bacterial species or populations is posed by genetic recombination. This is because there is a possibility that a single locus representing a particular strain may have undergone a recombination event with another strain or species and this locus would not be representative of the “true” evolutionary pathways of that particular strain genome. In other words, the use of a single locus will infer the evolution of this particular locus but not necessarily the evolution of the organism as a whole. MLST schemes aim at overcoming this problem by combining several, often seven, loci that are scattered across the genome. Thus, if one region of the genome has undergone recombination only one or two of the seven genes may be affected. This means primarily that if recombination is occurring it is easier to identify it by comparing base pair changes in the loci of closely related strains or the linkage between genes (Didelot and Falush, 2007; Feil et al., 2000). Secondly, in MLST schemes each allele of each gene is given a unique number so that isolates can be characterised by a multi-integer number called an allelic profile. This means that, regardless of whether a particular strain differs from another strain in a single locus by a single base pair (indicative of mutation) or many base pairs (indicative of recombination), in terms of the allelic profile, the strains will only differ by a single integer number. Thus analysing strains using their allelic profiles will buffer the distorted effect recombination may have on phylogenetic inferences or any other analyses.

Once the genes have been selected and the MLST scheme is in place, sequence data, strain information and allelic profiles are compiled by “virtual isolate collections centres” in the form of online databases (Urwin and Maiden, 2003) such as

1 www.mlst.net (Aanensen and Spratt, 2005). Each unique allelic profile is given a
2 unique number called a sequence type (ST) allowing for easy reference to particular
3 isolates. The original aim of the MLST concept was to enhance clinical diagnosis,
4 epidemiological monitoring, and population studies (Urwin and Maiden, 2003) but the
5 MLST concept has since been broadened to include the analysis of closely related
6 species and this approach has been named multi-locus sequence analysis (MLSA)
7 (Gevers et al., 2005; Hanage et al., 2006; Hanage et al., 2005). MLSA was developed
8 with the aim of allowing for rapid and robust hierarchical classification of all
9 prokaryotic species (Gevers et al., 2005) and has been raised as a solution to the time
10 consuming and complicated method of prokaryote species definition by DNA-DNA
11 hybridization (Bishop et al., 2009; Gevers et al., 2005). Recently a website has been
12 developed to allow the species identification of unknown isolates of Streptococcal
13 species (thought to be a taxonomically challenging group) by entering the sequence
14 data of seven gene fragments (Bishop et al., 2009).

15 For the LB group of spirochetes five schemes using multiple loci have been
16 developed (Table 2) (Bunikis et al., 2004; Margos et al., 2008; Qiu et al., 2004;
17 Richter et al., 2006; Rudenko et al., 2009a) and recently a mixture of two typing
18 schemes was used (Gomez-Diaz et al., 2011). Three of these schemes have been used
19 as an alternative to DNA-DNA hybridization, i.e. to delineate new species (Chu et al.,
20 2008; Margos et al., 2010; Margos et al., 2009; Postic et al., 2007; Richter et al.,
21 2006; Rudenko et al., 2009a; Rudenko et al., 2009b). Schemes by Bunikis et al.
22 (2004), Qiu et al. (2004) and Rudenko et al. (2009a) have tended to focus on species
23 found in the United States, with two focusing almost entirely on *B. burgdorferi* (Attie
24 et al., 2007; Brisson et al., 2010; Bunikis et al., 2004; Qiu et al., 2004). However,
25 most of these schemes did not adhere to the strict criteria set out by Urwin and

Maiden (2003), described above, because they combine a variety of gene types including slowly evolving housekeeping genes, non-coding regions, or fast evolving plasmid encoded loci. The loci differ in terms of the selective processes acting upon them, the number of variable sites within these loci as well as the gene category. This may lead to problems when inferring phylogenies as combining sequence data that are heterogeneous, as loci of different functional categories frequently are, can reduce the power of phylogenetic inference algorithms or even produce erroneous phylogenies (Huelsenbeck et al., 1996). Furthermore, the use of the 5S-23S IGS region as well as *ospA* means there is no species available to act as an outgroup to root a phylogeny and to allow for evolutionary inferences. For the MLSA scheme based on housekeeping genes (Margos et al., 2008), a website (borrelia.mlst.net) is maintained at Imperial College London, UK. It currently contains data for approximately 1,200 *Borrelia* strains comprising most of the described LB group species which have been resolved into >300 STs from Europe, Asia, and North America. The accumulative nature of MLST databases and the additional information gathered (e.g. geographic coordinates) makes it an attractive instrument to understand intra- and inter-species relationships on a global and regional scale.

4. *Borrelia* taxonomy

Bacterial taxonomy is a scientific discipline in flux (Gevers et al., 2006). For many years in bacterial systematics the accepted species definition was that a species would include strains with greater than 70 % homology when tested by DNA-DNA hybridization and a ΔT_m of 5°C or less. Below the value of 70 % homology strains were considered different species (Wayne et al., 1987). DNA-DNA hybridization requires a specialized laboratory and the number of laboratories that can perform this

analysis worldwide is limited. There are also questions about the interpretation and reproducibility of the method (Stackebrandt and Ebers, 2006). As this method is complicated, sequencing of the 16S rRNA locus and phylogenetic analysis was a valuable and widely used tool for bacterial classification. Both these methods, however, lacked sensitivity at the species level (Staley, 2006). Multilocus sequence analysis (MLSA), the genus-wide application of MLST, was proposed as an alternative to DNA-DNA hybridization and this technique is increasingly used in bacterial classification (Bishop et al., 2009; Gevers et al., 2006).

For LB group spirochetes, in addition to DNA-DNA hybridization and 16S sequences, analyses of the 5S-23S IGS have also served for species and strain typing (Postic et al., 1994). Several species have been defined using these methods including *B. burgdorferi* B31, *B. afzelii* VS461, *B. garinii* 20047, *B. japonica* HO14, *B. valaisiana* VS116 and *B. lusitaniae* PotiB2 (Baranton et al., 1992; Johnson et al., 1984; Kawabata et al., 1993; Le Fleche et al., 1997; Wang et al., 1997). In MLSA analyses these LB species cluster monophyletically at the end of long branches separating the different species (Margos et al., 2009; Richter et al., 2006) (see Figure 2 A).

For *Borrelia* taxonomy, the different schemes using multiple loci employed varying loci (Table 2). These schemes have been used to define several new *Borrelia* species (i.e. *B. spielmanii*, *B. californensis*, *B. carolinensis*, *B. americana*, *B. yangtze*, *B. bavariensis* and *B. kurtenbachii*) by genetic distance analyses (Chu et al., 2008; Margos et al., 2010; Margos et al., 2009; Postic et al., 2007; Richter et al., 2006; Rudenko et al., 2009a; Rudenko et al., 2009b). Richter and colleagues (Richter et al., 2006) and Postic and colleagues (Postic et al., 2007) compared the genetic distances of strains, based on the concatenated sequence of multiple genes, to the corresponding

whole DNA-DNA hybridization genetic distance data and determined a ‘cut-off’ value for species determination for their scheme (Postic et al., 2007; Richter et al., 2006). To determine this cut-off value, two European *B. burgdorferi* strains, NE49 and Z41293, which were ‘borderline’ *B. burgdorferi* stains in DNA-DNA hybridization, were used (Postic et al., 2007). A recently proposed 19th species, *B. finlandensis* (Casjens et al., 2011), belongs to this group of ‘borderline’ *B. burgdorferi* strains as determined by MLSA (see borrelia.mlst.net). While for the Richter-scheme the cut-off value was determined to be 0.21, using the same strains, Margos and co-authors (Margos et al., 2009) determined a cut-off value of 0.170 for the scheme based on eight chromosomally located housekeeping genes. This scheme permitted *B. bavariensis*, a rodent-associated ecotype previously named *B. garinii* OspA serotype 4, to be distinguished from other bird-associated *B. garinii* strains. This MLSA system enabled Takano and co-authors to determine that in Japan most human-pathogenic *Borrelia* isolates were phylogenetically closer related to the rodent-adapted sequence types ST84 and ST85 (*B. bavariensis*) than to bird-associated *B. garinii* (Takano et al., 2011).

5. Geographic distribution

The LB species are not evenly distributed across the globe (Figure 4). Host specialization and/or vector compatibility of the LB spirochetes are likely to influence the global distribution of different spirochetal species. In Europe, eight species have been recorded of which three (*B. garinii*, *B. afzelii*, and *B. bavariensis*) are also found throughout Asia (Baranton et al., 1992; Korenberg et al., 2002; Masuzawa, 2004; Takano et al., 2011). *B. valaisiana*, which occurs sympatrically with *B. garinii* in Europe, has rarely been found in *I. persulcatus* and appears to be absent in Russia and

1 most of Asia except for a single strain that was found in *I. columnae* in Japan
2 (Bormane et al., 2004; Korenberg et al., 2002; Masuzawa, 2004). Similarly, *B.*
3 *burgdorferi* has not been found in *I. persulcatus*, a main vector of *B. afzelii*, *B. garinii*
4 and *B. bavariensis*-like strains in Russia and Asia. Furthermore, NT29 strains of *B.*
5 *garinii* (which are rodent-adapted and genetically closely related to *B. bavariensis*
6 (unpublished)) occur in Russia and Asia but have not been found in *I. ricinus*
7 (Korenberg et al., 2002; Masuzawa et al., 2005). These authors concluded that the
8 distribution range of NT29 strains is associated with that of a single vector species, *I.*
9 *persulcatus*. This is interesting in view of the close phylogenetic relationship that has
10 been found for *B. bavariensis* (which is transmitted by *I. ricinus*) and rodent-adapted
11 *B. garinii* from Asia (Takano et al., 2011) and could provide an attractive system to
12 investigate *Ixodes* vector adaptations of *Borrelia* species. Species with a localised
13 distribution are *B. tanukii*, *B. turdi*, and *B. japonica* in Japan (Fukunaga et al., 1996b)
14 and *B. lusitaniae* around the Mediterranean Basin. Lizards of the family Lacertidae
15 have been identified as important hosts for the latter species (Amore et al., 2007;
16 Richter and Matuschka, 2006; Younsi et al., 2005). Whether or not other hosts are
17 reservoir competent has not been shown but, occasionally, infections of questing ticks
18 with *B. lusitaniae* have been described in other parts of Europe such as Poland and
19 Latvia (Vollmer et al., 2010; Wodecka and Skotarczak, 2005). *B. garinii* possibly has
20 the broadest distribution of all the LB group spirochetes. Not only is it found in
21 forested regions across Eurasia, it is also maintained in sea bird colonies by the tick
22 vector, *I. uriae*. This means it is also found in many far reaching sites including arctic
23 regions and colonies off the east coast of Canada (Duneau et al., 2008; Smith et al.,
24 2006). At first sight it is surprising, given the wide distribution of *B. garinii* and the
25 apparent overlap of terrestrial and seabird cycles in Europe (Comstedt et al., 2006)

1 that in North America, *B. garinii* has not spread into inland areas and remains limited
2 to coastal regions of Newfoundland (Smith et al., 2006). However, the lack of tick
3 vectors that could maintain terrestrial transmission cycles in this region is likely a
4 major reason why the seabird cycles have not spilled over into the rest of North
5 America (Ogden et al., 2009a).

6 Differences in *Borrelia* transmission cycles also exist at a much finer
7 scale driven by ecological factors, habitat types and microclimate which may locally
8 determine tick and host abundance (Eisen et al., 2006; Fingerle et al., 2004; Hubalek
9 and Halouzka, 1997; Killilea et al., 2008; Piesman, 2002; Rauter and Hartung, 2005).

10 Of the named species, seven occur in North America including *B. andersoni*,
11 *B. bissettii*, *B. californensis*, *B. carolinensis*, *B. americana*, *B. kurtenbachii* and *B.*
12 *burgdorferi* (Figure 4). *B. bissettii* has been found in Colorado, Illinois, California,
13 North Carolina and South Carolina where *I. spinipalpis*, *I. pacificus*, or *I. affinis* act as
14 vectors (Bissett and Hill, 1987; Lin et al., 2003; Maggi et al., 2010; Maupin et al.,
15 1994; Norris et al., 1999; Picken et al., 1995; Postic et al., 1998). Although it had
16 been reported that *B. bissettii* can be transmitted by *I. scapularis* under experimental
17 conditions (Oliver, 1996), *B. bissettii* has not been found in questing *I. scapularis*.
18 Similarly, *B. kurtenbachii* has been isolated from host-derived larvae and DNA has
19 been isolated from one questing adult *I. scapularis* (Anderson et al., 1988; Ogden et
20 al., 2011; Picken and Picken, 2000) but the species has rarely been found in *I.*
21 *scapularis* dominated habitats in recent years (Gatewood et al., 2009; Hamer et al.,
22 2007; Hoen et al., 2009; Ogden et al., 2011; Oliver et al., 2006). If generalist vectors
23 are able to transmit under experimental conditions LB group species that are usually
24 transmitted by endophilic vectors, the question arises, why does it not happen more

frequently in natural transmission cycles, why are these species not more widely distributed and what limits their distribution?

The species with the widest distribution in North America is *B. burgdorferi*, ranging from NE, to Upper Midwest (MW) and Western States. It also occurs in some Southern States and Southern Canada, within the distribution ranges of *I. scapularis*, *I. pacificus*, and *I. affinis*. Interestingly, in North Carolina, it was found predominantly in *I. affinis* but not *I. scapularis* and occurred sympatrically with *B. bissettii* (Maggi et al. 2010). In the NE *B. burgdorferi* appears to be the only LB species transmitted by *I. scapularis*. Climatic conditions impacting tick phenology may favour selection of certain strains (Diuk-Wasser et al., 2006; Gatewood et al., 2009; Ogden et al., 2007). Both, *B. burgdorferi* and *B. bissettii* have been recorded in Europe and North America (Postic et al. 1998, Gern and Humair 2002). In Europe, *B. bissettii* has been mainly described from human patients (Picken et al., 1996a; Picken et al., 1996b; Rudenko et al., 2008) but has only rarely been found in questing *I. ricinus* ticks (Hulinska et al., 2007). Curiously, human infection with *B. bissetti* in the USA has not been reported. Continued study of field-collected samples will likely continue to increase the number of known *Borrelia* species (Scott et al., 2010).

6. Population structure and dispersal patterns of LB species

Host specialization is an important factor in vector-borne disease, and different vector-borne pathogens show varying levels and patterns of host specialization. An accurate understanding of the epidemiology of many zoonoses can only be achieved by considering the varied ecological adaptations of the pathogens, particularly differences in host specificity (Dubska et al., 2009; Hanincova et al., 2003a; Hanincova et al., 2003b; Hu et al., 1997; Hu et al., 2001; Huegli et al., 2002;

Kurtenbach et al., 2001; Taragel'ova et al., 2008). The variation in host specialization makes the LB group of spirochetes an ideal model to directly contrast the effects of host specialization on the distribution of pathogens. As ticks cannot move over large distances independently (Falco and Fish, 1991), it has been suggested that the spread of LB spirochetes is linked to the movement of their hosts (Kurtenbach et al., 2002). In addition to being of public health importance, the delineation and monitoring of the geographic ranges of the different LB species also provides opportunities to examine in more general terms the role of host ecology in the epidemiology of vector-borne zoonoses.

6.1 Europe and Asia

MLSA on housekeeping genes has revealed differences in the level of geographic structuring of populations of LB species that are consistent with distribution patterns of their different vertebrate hosts (Vitorino et al., 2008; Vollmer et al., 2011).

Vitorino and co-authors (2008) investigated *B. lusitaniae*, a species that has been associated with lizards, from two geographic regions in Portugal (Mafra and Grandola), which are approximately 160 km apart and located north and south of Lisbon. A pronounced fine-scale phylogeographic population structure was observed where most strains from Mafra clustered separately from Grandola strains (Vitorino et al., 2008). The authors suggested that this distribution reflects the highly parapatric population structure of lizards on the Iberian peninsula (Paulo et al., 2008).

Vollmer and co-authors (Vollmer et al., 2011) tested the prediction that host movement determines spirochaete biogeography by characterising *B. garinii*, *B. valaisiana*, and *B. afzelii* from various sites in Europe (Great Britain, France, Germany, Latvia). MLSA of the rodent-associated species, *B. afzelii*, showed a

1 population structure that signified restricted movement of strains between geographic
2 regions. This differentiation was pronounced: only two *B. afzelii* STs have been found
3 in more than one geographic location (Figure 5, Panel C). These data suggested that
4 the English Channel may act as a barrier to the movement of *B. afzelii* strains between
5 Great Britain and continental Europe (Vollmer et al., 2011). Chinese and European *B.*
6 *afzelii* populations also showed high levels of differentiation suggesting very limited
7 movement over these large distances. However, one Chinese *B. afzelii* strain clustered
8 within the European group suggesting that there may be rare cases of movement
9 between East and West, although the mechanisms behind such events are unclear
10 (Vollmer, personal communication).

11 The data obtained by Vollmer and co-authors (Vollmer et al., 2011) are
12 suggestive of interesting parallels between *B. afzelii* and the evolutionary history of
13 their vertebrate host. In phylogenies and eBURST analyses, *B. afzelii* strains from
14 Scotland appeared to be more closely related to STs found in Latvia than to STs
15 found in England, suggesting that there is limited, or potentially no, movement of *B.*
16 *afzelii* between north and south in the UK. This is interesting in the light of studies
17 that investigated phylogenetic relationships of small mammals (including the field
18 vole *Microtus agrestis*, bank vole *Myodes glareolus*, and pygmy shrew *Sorex*
19 *minutus*) in Great Britain which showed a clear north/south divide between
20 phylogroups (Searle et al., 2009). The marked differentiation between English and
21 Scottish *B. afzelii* samples may therefore be a result of limited north-south rodent
22 dispersal, although this hypothesis needs further investigation (Vollmer et al., 2011).
23 In addition, other studies of potential host species of *B. afzelii* including shrew and
24 vole species have observed phylogeographic structuring of populations across Europe.
25 These studies have attributed the phylogeographic patterns to population expansions

1 from ancestral refugia after the last glacial maximum (LGM), which possibly included
2 an Iberian and an East Baltic refuge (Heckel et al., 2005; Hewitt, 1999, 2001; Taberlet
3 and Bouvet, 1994; Taberlet et al., 1998). Northward spread of the populations from
4 the two refugia led to a possible overlap in the region of Germany or the Czech
5 Republic (Figure 6) (Heckel et al., 2005; Hewitt, 1999). Data of European *B. afzelii*
6 strains bear some resemblance of populations maintained potentially by the two
7 mammalian refuge populations as the *B. afzelii* phylogeny could be divided into a
8 Western European cluster and an Eastern European cluster (Vollmer, personal
9 communication).

10 However, fine scale structuring can also be observed in vole species either due
11 to natural and man made barriers (e.g. large rivers, highways) or due to a social
12 structure within population. These processes may limit the rates of movement
13 between host populations (Gerlach and Musolf, 2000; Schweizer et al., 2007) and
14 therefore limit dispersal of *B. afzelii*. Observations of *B. afzelii* strains at one site in
15 Latvia and the English sites are consistent with fine-scale structuring of the bacterial
16 populations due to restricted host movements (Vollmer et al., 2011). However, *B.*
17 *afzelii* has many rodent host species and further studies of small mammal host species
18 of *B. afzelii* may be required to better understand the ability of this species to disperse.

19 In contrast, both of the bird-related species investigated, *B. valaisiana*
20 and *B. garinii*, showed evidence of spatial mixing of STs between geographic regions
21 (Figure 5A, B) (Vollmer et al., 2011). Interestingly, while *B. garinii* data suggested
22 free movement of strains, *B. valaisiana* showed low to moderate differentiation,
23 suggesting there is not complete homogenization of *B. valaisiana* strains within
24 Europe. This was surprising because both species have been reported to be maintained
25 by similar species of avian hosts (Dubska et al., 2009; Taragel'ova et al., 2008) but

may suggest subtle ecological differences between these species. Certainly *B. garinii* differs from *B. valaisiana* in being maintained in cycles between seabirds and their associated tick, *I. uriae* (Bunikis et al., 1996; Larsson et al., 2007; Olsen et al., 1995; Olsen et al., 1993) as well as in terrestrial cycles. Several studies (Comstedt et al., 2006; Gomez-Diaz et al., 2011) reported an overlap of marine and terrestrial *B. garinii* populations but the full impact on the observed population structure remains to be investigated. Notably, *B. garinii* STs from China showed divergence from European *B. garinii* STs indicated by long branches joining them to their closest European relatives in phylogenetic inferences (Vollmer, personal communication) and suggesting limited gene flow between the two regions. These data also suggested that the role that migratory birds play in east-west or west-east movement of *B. garinii* may be limited as would be expected as most migratory bird movement is on the north-south axis. Analyses of more Russian and Asian *B. garinii* samples would be required to confirm this hypothesis and assess the level of movement of *B. garinii* between Asia and Europe.

Given that the movement of some LB species is limited by the propensity for their vertebrate hosts' ranges to shift, landscape genetic analysis would be an appropriate approach to determine barriers to movement (Manel et al., 2003). Such future investigations will be facilitated by identifying the full host spectrum of the different LB species.

6.2 North America

In North America a complex picture of LB group species has emerged. While habitats in California and the Southeastern States harbour a great variety of LB species, the prevalence of human infections is low (Bacon et al., 2008). This may be related to

1 host preferences and human biting behaviour of main vectors in these regions or other
2 ecological factors, although it may also be due to some of these species being non-
3 pathogenic in humans (Eisen et al., 2004; Eisen et al., 2009; Girard et al., 2011; Lane
4 and Quistad, 1998; Norris et al., 1996; Oliver, 1996; Oliver et al., 2003; Piesman,
5 1993; Swei et al., 2011; Talleklint-Eisen and Eisen, 1999; Wright et al., 1998). In
6 Southeastern States, infection prevalence in *I. scapularis* ticks is low, which may be
7 due to ‘dilution’ of transmission cycles by reservoir-incompetent lizards acting as tick
8 hosts, and or by climate-driven tick seasonality that is less favourable for transmission
9 (Durden et al., 2002; Ogden et al., 2008a; Ogden and Tsao, 2009b; Spielman et al.,
10 1984); (Kollars et al., 1999; Spielman et al., 1984; Swanson and Norris, 2007).

11 In the NE USA, *B. burgdorferi* is the predominant species, human infection
12 incidence is high (>80 % of all recorded infections in the USA occur here), and, here
13 the first population level studies on *B. burgdorferi* were conducted. Pioneering studies
14 using *ospA* and *ospC* as genetic markers (Qiu et al., 1997; Qiu et al., 2002; Wang et
15 al., 1999) found a high local variation of strains but a uniform distribution across the
16 NE USA. The authors suggested that ancient polymorphisms combined with
17 balancing selection (in form of negative frequency-dependent immune selection)
18 maintains the high diversity of populations (Dykhuizen et al., 1993; Qiu et al., 2002;
19 Wang et al., 1999). Parallel studies on *I. scapularis* populations suggested that
20 migration could also be at play as ‘American clade’ *I. scapularis* (Norris et al., 1996)
21 were found in coastal bird sanctuaries in North Carolina (Qiu et al., 2002). Further
22 studies on *ospC* including European and American strains of *B. burgdorferi* led the
23 authors to suggest that recent and rapid spread of *B. burgdorferi* across two continents
24 has occurred (Qiu et al. 2008). Transportation of ticks by infected migratory birds has
25 more recently been suggested for the introduction of *B. burgdorferi* strains and *I.*

1 *scapularis* ticks into southern Canada (Ogden et al., 2010; Ogden et al., 2008b; Ogden
2 et al., 2011; Ogden et al., 2006). Although both balancing selection and migration
3 may have a homogenizing effect, there are several lines of evidence supporting the
4 argument for balancing selection and/or functional constraints (related to its role in
5 tissue adherence or protein binding during invasion/infection processes) acting on
6 *ospC*: 1) identical *ospC* major types are found in all *B. burgdorferi* populations but
7 these are regionally matched with different MLST STs (Margos et al., 2008; Qiu et
8 al., 2008; Travinsky et al., 2010). This suggests that slowly evolving housekeeping
9 genes have accumulated mutations while the *ospC* gene has not. 2) The description of
10 *ospC* types that are found exclusively in Europe (e.g. P, Q, S, V) or California (e.g.
11 H3, E3) points to population separation as frequent exchange between the populations
12 would homogenize alleles (Girard et al., 2009; Qiu et al., 2008; Wang et al., 1999).
13 The finding of these ‘private’ *ospC* types has been interpreted as adaptation to new
14 habitats (Girard et al., 2009; Qiu et al., 2008). It could - alternatively - reflect loss of
15 *ospC* major types in some regions due to severe population bottlenecks as described
16 for the USA (Spielman, 1994).

17 MLST data based on housekeeping genes paint a different picture for *B.*
18 *burgdorferi* populations. These data support the view that the *B. burgdorferi*
19 populations from Europe and North America and in North America are genetically
20 related but are currently separated with no or limited gene flow between them (Hoen
21 et al., 2009; Margos et al., 2008; Ogden et al., 2011). In 2004 a CDC project was
22 launched to investigate the presence and infection prevalences of *I. scapularis*
23 nymphs on a country-wide scale and questing nymphs were collected systematically
24 from May to September (Diuk-Wasser et al., 2006). Hoen and co-authors (Hoen et al.,
25 2009) investigated the population structure of *B. burgdorferi* by MLST using 78

1 samples from 2004 and 2005: 41 samples were from NE sites and 37 from MW sites.
2 Thirty seven distinct STs were determined but no single ST was found in both regions
3 suggesting restricted present day gene flow between the two regions. It further
4 suggested that the coincident emergence of Lyme borreliosis in the two regions
5 originated from multiple expansions of vector tick and *B. burgdorferi* populations
6 (Hoen et al., 2009). Although the observed level of sequence divergence in some
7 samples from NE and MW was only few nucleotides, considering the slow evolution
8 of housekeeping genes, these mutational changes may have accumulated over time
9 periods that exceeded the latest Lyme disease emergence in North America in the past
10 40 years. These results were supported by additional studies suggesting limited gene
11 flow between the *B. burgdorferi* populations described from the NE and Upper MW
12 ((Ogden et al., 2011), Bent, personal communication).

13 In a follow-up study, two transects in the NE were intensely sampled for
14 questing *I. scapularis* nymphs in 2007, one transect starting from Long Island
15 following the Hudson River valley north to Lake Champlain in Vermont, the other
16 one starting in Old Lyme going north into Massachusetts. Although most STs were
17 found in all sites, regression analysis revealed differences in frequencies of ST which
18 correlated with latitude (Bent, personal communication). The mechanism behind the
19 observed structure is currently unknown but possible reasons could be differences in
20 host composition, pattern of movement, or genetic drift. Genetic drift can prevail and
21 weaken natural selection in populations with small effective population size (N_e)
22 ((Page and Holmes, 1998). Due to their parasitic life style, LB spirochetes are thought
23 to have a small N_e (Qiu et al., 2002). Interestingly, although more than 300 samples
24 were analysed, the number of new STs was small: four new STs were found in
25 addition to the 37 STs described by Hoen et al. (Hoen et al., 2009) from the NE.

1 These data are consistent with limited genetic diversity in this region of high LB
2 incidence probably reflecting the severe bottleneck that has been suggested previously
3 (Spielman, 1994). Similar studies on *I. scapularis* samples collected by passive
4 surveillance in Canada (from Nova Scotia to Manitoba) also supported the notion of
5 geographic separation with limited gene flow between NE and MW. STs determined
6 east of 80° longitude resembled those of NE USA, while STs west of 80° longitude
7 resembled those of MW *B. burgdorferi* populations. Geographic analysis of STs and
8 *ospC* alleles were consistent with south-to-north dispersion of infected ticks from the
9 USA, likely on migratory birds (Ogden et al., 2011). Surprisingly, 19 novel STs were
10 determined which were single (SLV), double (DLV) or triple locus variants (TLV) of
11 STs from the USA supporting the notion that the spatial scale of sampling is
12 important to capture the population variation of, and to understand demographic
13 processes in, *B. burgdorferi* (Figure 7). Preliminary MLST data from approximately
14 25 strains from the Upper MW and 25 strains from California show that additional
15 samples from these regions led to denser eBurst ‘forests’ and better resolution of
16 clonal complexes. Indeed, in the Californian dataset the first SLV of ST1 (B31) was
17 determined (Margos, unpublished) supporting the view that *B. burgdorferi*
18 populations across North America – not only in the MW and NE but also from
19 California – are genetically related and once belonged to an admixed population
20 (Hoen et al., 2009).

21 The complexity observed for *B. burgdorferi* populations in North America is
22 likely due to a dynamic short- and long-term evolution. The long-term evolutionary
23 history was probably shaped by glacial-interglacial cycles (Humphrey et al., 2010;
24 Qiu et al., 2002) which is consistent with data by Hoen et al. (2009) who found
25 signatures of ancient population expansions of *B. burgdorferi* likely to date back

1 several thousand, if not millions of years ago. Demographic events in the past 200
2 years (following the arrival of European settlers) have shaped populations of hosts
3 and vectors by deforestation, dwindling deer and tick populations and causing severe
4 bottlenecks in *Borrelia* populations (McCabe and McCabe, 1997; Spielman, 1994).
5 Since then, expansion of deer and tick populations have resulted in the latest dispersal
6 of LB spirochetes leading to an epidemic of human LB in the NE and MW USA
7 during the past four decades (Bacon et al., 2008). It is conceivable that different
8 regions were affected in different ways by these processes but in order to understand
9 the contemporary pattern sequence data with high resolution power, such as genome
10 wide SNPs, will be required (Figure 9).

11 *B. burgdorferi* is considered a generalist species that can be maintained and
12 transmitted to ticks by a great variety of hosts including birds and rodents (Brisson
13 and Dykhuizen, 2006; Hanincova et al., 2006; Richter et al., 2000), therefore,
14 understanding its dispersal is more complicated than that of host specialized species.
15 Fitness variation in hosts has been described for several strains (Derdakova et al.,
16 2004; Hanincova et al., 2008b) and host adaptations may be developing (Brinkerhoff
17 et al., 2010; Brisson and Dykhuizen, 2004; Ogden et al., 2011); all of which is likely
18 to impact transmission efficiency and dispersal of *B. burgdorferi* strains. Some
19 models of dispersal of *B. burgdorferi* have emerged that are consistent with slow
20 south-to-north range expansions of *B. burgdorferi* that lag behind expansion of the
21 tick vector (Ogden et al., 2010). Whether or not there is low level east-west or west-
22 east migration, is far less understood (Hamer et al., 2010; Ogden et al., 2011). Clearly,
23 additional information on the ecology of *B. burgdorferi* strains is required in order to
24 obtain a comprehensive picture of how *B. burgdorferi* strains spread.

25

7. Models of global evolution

For phylogenetic analyses of the whole group of LB spirochetes, housekeeping genes provide the benefit of defining outgroup species as they are also present in the relapsing fever spirochetes (*B. hermsii*, *B. duttoni*, *B. turicatae*) allowing rooting of phylogenies. Their analysis also allows inferences of the temporal evolution of LB species. However, ascertaining this order using a single gene such as *flaB*, or even MLSA, proved difficult due to low confidence values of internal branches in phylogenies in which all STs of the European species were included (Fukunaga et al., 1996c; Kurtenbach et al., 2010). Several factors may be responsible for this: 1) internal branches representing species divisions are extremely short suggesting that the speciation events, in evolutionary terms, occurred in quick succession. Thus there are limited mutations existing in the sequences today that represent these intermediate species. 2) The limited number of genes may not contain a sufficient number of nucleotide polymorphisms to clearly define the topology. 3) These short branches may be suggestive of incomplete lineage sorting (Avice and Robinson, 2008; Maddison and Knowles, 2006). This occurs when polymorphisms are maintained in a gene through two or more speciation events, thus giving the impression of a different topology.

Several unrooted or midpoint rooted phylogenies have been published for LB group species (Margos et al., 2010; Richter et al., 2006; Rudenko et al., 2009b) (Figure 8). These trees produced different topologies compared to each other and to the concatenated housekeeping gene trees. Differences in tree topology are most notable comparing phylogenetic inferences for MLSA genes and *ospC* suggesting different evolutionary pathways of plasmid encoded and chromosomal genes (Figures 2A, B).

1 There are, however, some species that form clusters in all trees generated
2 using single or multiple chromosomal loci. Notably, the ‘American’ species and the
3 ‘Eurasian’ species form sister clades joined by a well supported branch suggesting
4 that these two clades separated early during LB evolution. Within the ‘American’
5 clade, *B. burgdorferi* and *B. bissettii* (both occurring in North America and Europe)
6 fall into different subclades raising questions about migration times and routes
7 between continents. Several species (if included in phylogenies) tend to always cluster
8 closely together such as *B. afzelii* and *B. spielmanii* or *B. garinii* and *B. bavariensis*
9 being consistent with more recent speciation events (Margos et al., 2010; Postic et al.,
10 2007; Rudenko et al., 2009b). It is also apparent that host associations did not develop
11 only once. For example, not all bird-adapted LB species cluster monophyletically in
12 the species tree (Figure 2A) suggesting that several host switches occurred during the
13 evolutionary history of LB species.

14 The doubling time of LB spirochetes in feeding nymphs has been estimated to
15 be four hours but was much slower *in vitro*, approximating 8-12 h under constant
16 temperature conditions (33°C) (De Silva and Fikrig, 1995; Heroldova et al., 1998;
17 Pollack et al., 1993) and may be even longer at lower (winter) temperatures in vector
18 populations under natural conditions. It is, therefore, extremely difficult to estimate
19 mutation rates or time of speciation events for LB species by comparison with other
20 bacterial species. Consequently, to establish a realistic time frame of LB species
21 evolution, measures of mutation rates for LB species are essential which will require
22 the use of larger sets of sequence data than MLSA.

24 **8. Future Avenues**

1 In this paper we have summarized recent research on population genetics, molecular
2 taxonomy and evolution of LB spirochetes which has moved from single locus
3 approaches to multilocus approaches. From the information gathered here, it is
4 evident that major advances have been made in understanding the evolutionary
5 ecology of LB spirochetes but there are also limitations which need to be addressed.
6 These include questions like: 1) What drives associations/adaptation between LB
7 group spirochetes and their hosts and vectors? 2) What is the full host spectrum of the
8 different LB species? 3) Which factors apart from host associations impact dispersal
9 of LB group species? 4) What is the speed and geometry of spread?

10 Some methods have been developed to address such questions. For example
11 blood meal analyses analysis in questing ticks may help to resolve host associations
12 (Humair et al. 1997) and real time genotyping assays are already being used for LB
13 spirochetes for loci such as IGS, *fla* or *hbb* (Herrmann and Gern, 2010; Portnoi et al.,
14 2006; Strube et al., 2010), but these methods may need refinement. Next generation
15 sequencing and SNP analyses will likely prove very valuable to develop better tools
16 for precise strain identification, to identify mixed infections in ticks or patients, to
17 refine blood meal analysis or to address questions regarding the deep evolutionary
18 relationships of LB group spirochetes. Developments such as single nucleotide primer
19 extension assays (Murphy et al., 2003) or high melting resolution techniques (Wittwer
20 et al., 2003) may be suitable for such approaches.

21 MLST of housekeeping genes in LB spirochetes has shown that recombination
22 can occur on chromosomally located loci. In general, the use of MLST/MLSA has
23 shown that there is great variation in bacterial inheritance. While some taxa such as
24 *Staphylococcus aureus*, *Yersinia* sp or *Salmonella typhi* show little horizontal gene
25 transfer, others show an enormous amount of recombination or horizontal gene

transfer, well known examples are *Neisseria* sp and *Helicobacter pylori* (Achtman, 2004; Feil et al., 2003; Feil and Spratt, 2001; Holt et al., 2008; Ochman et al., 2000). However, the availability of whole genome sequences for a large numbers of closely related bacteria has led to the concept that most bacterial genomes consists of a ‘core’ genome which can inform about evolutionary relationships and an ‘accessory’ genome which is much more flexible, permits invasion of new niches or confers selective advantages (such as antibiotic resistance) by horizontally acquired genome elements (often plasmids) (Guttman and Stavrinides, 2010; Ochman et al., 2000).

For some microbial pathogens whole genome sequencing and detection of SNPs has led to a quantum leap forward in understanding evolutionary and demographic processes. Fortunately, the haploid nature of bacteria makes it convenient to detect these processes using sequence data. Genome-wide SNPs permit us to distinguish evolutionary processes, such as mutation, recombination, selection, and drift, from demographic processes affecting the whole genome such as migration, population expansion/contractions (Guttman and Stavrinides, 2010). Analyses of genome-wide SNPs permit much more accurate estimates of mutation rates which is a vital parameter in population level and evolutionary studies, as well as the analyses of recent as well as ancient events (Figure 9). Computer programs have been developed for bacterial population based inferences and many are tailored for use with MLST/MLSA or SNP data (Corander and Marttinen, 2006; Corander et al., 2004; Didelot et al., 2009; Didelot and Falush, 2007; Feil et al., 2004; Francisco et al., 2009; Guillot, 2008; Guillot et al., 2008; Kuhner, 2006; Schierup and Wiuf, 2010) (see review by (Excoffier and Heckel, 2006)).

Several *Borrelia* genomes have been sequenced and for *B. burgdorferi* more than 10 draft genomes are available (Schutzer et al., 2011). While this is a good start

1 and can provide the scaffolding for next generation sequencing of further samples,
2 understanding the most recent population expansion in northeastern America requires
3 the analyses of carefully selected samples from that region. As MLST and eBurst data
4 convincingly demonstrate, getting insights into the deep evolutionary history of *B.*
5 *burgdorferi* requires sampling at a different scale. In our opinion – the time is ripe to
6 take *Borrelia* research to the next step, and that is the emerging field of bacterial
7 population genomics (Guttman and Stavrinides, 2010) as this together with MLST
8 will provide a framework for epidemiological, clinical and ecological studies.

10 **Acknowledgements**

11 We would like to thank S. J. Bent, J. Tsao and R. S. Lane for sharing unpublished
12 data, numerous colleagues and the German Collection of Microorganisms and cell
13 cultures (DSMZ) for providing *Borrelia* DNA and a large number of tick collectors
14 for their efforts. The authors are grateful for financially support received by The
15 Wellcome Trust (grant no. 074322/Z/04/Z), Public Health Agency of Canada, NIH-
16 NIAID (grant nos. AR041511; 5R21AI065848-03); USDA-ARS Cooperative
17 Agreement; G. Harold and Leila Y. Mathers Charitable Foundation; US Centers for
18 Disease Control and Prevention.

20 **References**

- 21 Aanensen, D.M., Spratt, B.G., 2005. The multilocus sequence typing network:
22 mlst.net. *Nucleic Acids Res* 33, W728-733.
23 Achtman, M., 2004. Population structure of pathogenic bacteria revisited. *Int J Med*
24 *Microbiol* 294, 67-73.
25 Amore, G., Tomassone, L., Grego, E., Ragagli, C., Bertolotti, L., Nebbia, P., Rosati,
26 S., Mannelli, A., 2007. *Borrelia lusitaniae* in immature *Ixodes ricinus* (Acari:
27 Ixodidae) feeding on common wall lizards in Tuscany, central Italy. *J Med*
28 *Entomol* 44, 303-307.

- 1 Anderson, J.F., Magnarelli, L.A., McAninch, J.B., 1988. New *Borrelia burgdorferi*
2 antigenic variant isolated from *Ixodes dammini* from upstate New York. J Clin
3 Microbiol 26, 2209-2212.
- 4 Attie, O., Bruno, J.F., Xu, Y., Qiu, D., Luft, B.J., Qiu, W.G., 2007. Co-evolution of
5 the outer surface protein C gene (*ospC*) and intraspecific lineages of *Borrelia*
6 *burgdorferi* sensu stricto in the northeastern United States. Infect Genet Evol 7, 1-
7 12.
- 8 Avise, J.C., Robinson, T.J., 2008. Hemiplasy: a new term in the lexicon of
9 phylogenetics. Syst Biol 57, 503-507.
- 10 Bacon, R.M., Kugeler, K.J., Mead, P.S., 2008. Surveillance for Lyme disease--United
11 States, 1992-2006. MMWR Surveill Summ 57, 1-9.
- 12 Balashov, Y.S., 1972. Bloodsucking ticks (Ixodoidea) - vectors of diseases of man
13 and animals. Miscellaneous Publications of the Entomological Society of America
14 8, 163-376.
- 15 Baranton, G., Postic, D., Saint Girons, I., Boerlin, P., Piffaretti, J.C., Assous, M.,
16 Grimont, P.A., 1992. Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia*
17 *garinii* sp. nov., and group VS461 associated with Lyme borreliosis. Int J Syst
18 Bacteriol 42, 378-383.
- 19 Baranton, G., Seinost, G., Theodore, G., Postic, D., Dykhuizen, D., 2001. Distinct
20 levels of genetic diversity of *Borrelia burgdorferi* are associated with different
21 aspects of pathogenicity. Res Microbiol 152, 149-156.
- 22 Barbour, A., Garon, C.F., 1988. The genes encoding major surface proteins of
23 *Borrelia burgdorferi* are located on a plasmid. Annals of the New York Academie
24 of Sciences 539, 144-153.
- 25 Barbour, A.G., Bunikis, J., Travinsky, B., Hoen, A.G., Diuk-Wasser, M.A., Fish, D.,
26 Tsao, J.I., 2010. Niche Partitioning of *Borrelia burgdorferi* and *Borrelia*
27 *miyamotoi* in the same tick vector and mammalian reservoir species. American
28 Journal of Tropical Medicine and Hygiene 81, 1120-1131.
- 29 Barbour, A.G., Travinsky, B., 2010. Evolution and Distribution of the *ospC* Gene, a
30 Transferable Serotype Determinant of *Borrelia burgdorferi*. MBio 1.
- 31 Bishop, C.J., Aanensen, D.M., Jordan, G.E., Kilian, M., Hanage, W.P., Spratt, B.G.,
32 2009. Assigning strains to bacterial species via the internet. BMC Biol 7.
- 33 Bissett, M.L., Hill, W., 1987. Characterization of *Borrelia burgdorferi* strains isolated
34 from *Ixodes pacificus* ticks in California. J Clin Microbiol 25, 2296-2301.
- 35 Boerlin, P., Peter, O., Bretz, A.G., Postic, D., Baranton, G., Piffaretti, J.C., 1992.
36 Population genetic analysis of *Borrelia burgdorferi* isolates by multilocus enzyme
37 electrophoresis. Infect Immun 60, 1677-1683.
- 38 Bormane, A., Lucenko, I., Duks, A., Mavtchoutko, V., Ranka, R., Salmina, K.,
39 Baumanis, V., 2004. Vectors of tick-borne diseases and epidemiological situation
40 in Latvia in 1993-2002. Int J Med Microbiol 293 Suppl 37, 36-47.
- 41 Brinkerhoff, R.J., Folsom-O'Keefe, C.M., Tsao, K., Diuk-Wasser, M.A., 2010. Do
42 birds affect Lyme disease risk? Range expansion of the vector-borne pathogen
43 *Borrelia burgdorferi* Frontiers in Ecology and the Environment.
- 44 Brisson, D., Dykhuizen, D.E., 2004. *ospC* diversity in *Borrelia burgdorferi*: different
45 hosts are different niches. Genetics 168, 713-722.
- 46 Brisson, D., Dykhuizen, D.E., 2006. A modest model explains the distribution and
47 abundance of *Borrelia burgdorferi* strains. Am J Trop Med Hyg 74, 615-622.
- 48 Brisson, D., Vandermause, M.F., Meece, J.K., Reed, K.D., Dykhuizen, D.E., 2010.
49 Evolution of northeastern and midwestern *Borrelia burgdorferi*, United States.
50 Emerg Infect Dis 16, 911-917.

- 1 Bunikis, J., Garpmo, U., Tsao, J., Berglund, J., Fish, D., Barbour, A.G., 2004.
- 2 Sequence typing reveals extensive strain diversity of the Lyme borreliosis agents
- 3 *Borrelia burgdorferi* in North America and *Borrelia afzelii* in Europe.
- 4 Microbiology 150, 1741-1755.
- 5 Bunikis, J., Olsen, B., Fingerle, V., Bonnedahl, J., Wilske, B., Bergstrom, S., 1996.
- 6 Molecular polymorphism of the lyme disease agent *Borrelia garinii* in northern
- 7 Europe is influenced by a novel enzootic *Borrelia* focus in the North Atlantic. J
- 8 Clin Microbiol 34, 364-368.
- 9 Burgdorfer, W., Barbour, A.G., Hayes, S.F., Benach, J.L., Grunwaldt, E., Davis, J.P.,
- 10 1982. Lyme disease-a tick-borne spirochetosis? Science 216, 1317-1319.
- 11 Canica, M.M., Nato, F., du Merle, L., Mazie, J.C., Baranton, G., Postic, D., 1993.
- 12 Monoclonal antibodies for identification of *Borrelia afzelii* sp. nov. associated
- 13 with late cutaneous manifestations of Lyme borreliosis. Scand J Infect Dis 25,
- 14 441-448.
- 15 Casati, S., Bernasconi, M.V., Gern, L., Piffaretti, J.C., 2004. Diversity within *Borrelia*
- 16 *burgdorferi* sensu lato genospecies in Switzerland by *recA* gene sequence. FEMS
- 17 Microbiol Lett 238, 115-123.
- 18 Casjens, S.R., Fraser-Liggett, C.M., Mongodin, E.F., Qiu, W.G., Dunn, J.J., Luft,
- 19 B.J., Schutzer, S.E., 2011. Whole genome sequence of an unusual *Borrelia*
- 20 *burgdorferi* sensu lato isolate. J Bacteriol 193, 1489-1490.
- 21 Chu, C.Y., Liu, W., Jiang, B.G., Wang, D.M., Jiang, W.J., Zhao, Q.M., Zhang, P.H.,
- 22 Wang, Z.X., Tang, G.P., Yang, H., Cao, W.C., 2008. Novel genospecies of
- 23 *Borrelia burgdorferi* sensu lato from rodents and ticks in Southwestern China. J
- 24 Clin Microbiol 46, 3130-3133.
- 25 Collares-Pereira, M., Couceiro, S., Franca, I., Kurtenbach, K., Schafer, S.M.,
- 26 Vitorino, L., Goncalves, L., Baptista, S., Vieira, M.L., Cunha, C., 2004. First
- 27 isolation of *Borrelia lusitaniae* from a human patient. J Clin Microbiol 42, 1316-
- 28 1318.
- 29 Comstedt, P., Bergstrom, S., Olsen, B., Garpmo, U., Marjavaara, L., Mejlom, H.,
- 30 Barbour, A.G., Bunikis, J., 2006. Migratory passerine birds as reservoirs of Lyme
- 31 borreliosis in Europe. Emerg Infect Dis 12, 1087-1095.
- 32 Corander, J., Marttinen, P., 2006. Bayesian identification of admixture events using
- 33 multilocus molecular markers. Mol Ecol 15, 2833-2843.
- 34 Corander, J., Waldmann, P., Marttinen, P., Sillanpaa, M.J., 2004. BAPS 2: enhanced
- 35 possibilities for the analysis of genetic population structure. Bioinformatics 20,
- 36 2363-2369.
- 37 Daniel, M., Danielova, V., Kriz, B., Jirsa, A., Nozicka, J., 2003. Shift of the tick
- 38 *Ixodes ricinus* and tick-borne encephalitis to higher altitudes in central Europe.
- 39 Eur J Clin Microbiol Infect Dis 22, 327-328.
- 40 De Silva, A.M., Fikrig, E., 1995. Growth and migration of *Borrelia burgdorferi* in
- 41 *Ixodes* ticks during blood feeding. Am J Trop Med Hyg 53, 397-404.
- 42 Dennis, D.T., Hayes, E.B., 2002. Epidemiology of Lyme Borreliosis, in: Gray, J.S.,
- 43 Kahl, O., Lane, R.S., Stanek, G. (Eds.), Lyme Borreliosis: Biology of the
- 44 Infectious Agents and Epidemiology of Disease. CABI Publishing, Wallingford,
- 45 pp. 251-280.
- 46 Derdakova, M., Dudioak, V., Brei, B., Brownstein, J.S., Schwartz, I., Fish, D., 2004.
- 47 Interaction and transmission of two *Borrelia burgdorferi* sensu stricto strains in a
- 48 tick-rodent maintenance system. Appl Environ Microbiol 70, 6783-6788.
- 49 Didelot, X., Darling, A., Falush, D., 2009. Inferring genomic flux in bacteria. Genome
- 50 Res 19, 306-317.

- 1 Didelot, X., Falush, D., 2007. Inference of bacterial microevolution using multilocus
2 sequence data. *Genetics* 175, 1251-1266.
- 3 Diuk-Wasser, M.A., Gatewood, A.G., Cortinas, M.R., Yaremych-Hamer, S., Tsao, J.,
4 Kitron, U., Hickling, G., Brownstein, J.S., Walker, E., Piesman, J., Fish, D., 2006.
5 Spatiotemporal patterns of host-seeking *Ixodes scapularis* nymphs (Acari:
6 Ixodidae) in the United States. *J Med Entomol* 43, 166-176.
- 7 Diza, E., Papa, A., Vezyri, E., Tsounis, S., Milonas, I., Antoniadis, A., 2004. *Borrelia*
8 *valaisiana* in cerebrospinal fluid. *Emerg Infect Dis* 10, 1692-1693.
- 9 Dolan, M.C., Piesman, J., Mbow, M.L., Maupin, G.O., Peter, O., Brossard, M., Golde,
10 W.T., 1998. Vector competence of *Ixodes scapularis* and *Ixodes ricinus* (Acari:
11 Ixodidae) for three genospecies of *Borrelia burgdorferi*. *J Med Entomol* 35, 465-
12 470.
- 13 Dubska, L., Literak, I., Kocianova, E., Taragelova, V., Sychra, O., 2009. Differential
14 role of passerine birds in distribution of *Borrelia* spirochetes, based on data from
15 ticks collected from birds during the postbreeding migration period in Central
16 Europe. *Appl Environ Microbiol* 75, 596-602.
- 17 Duneau, D., Boulonier, T., Gomez-Diaz, E., Petersen, A., Tveraa, T., Barrett, R.T.,
18 McCoy, K.D., 2008. Prevalence and diversity of Lyme borreliosis bacteria in
19 marine birds. *Infect Genet Evol* 8, 352-359.
- 20 Durden, L.A., Oliver, J.H., Jr., Banks, C.W., Vogel, G.N., 2002. Parasitism of lizards
21 by immature stages of the blacklegged tick, *Ixodes scapularis* (Acari, Ixodidae).
22 *Exp Appl Acarol* 26, 257-266.
- 23 Dykhuizen, D.E., Baranton, G., 2001. The implications of a low rate of horizontal
24 transfer in *Borrelia*. *Trends Microbiol* 9, 344-350.
- 25 Dykhuizen, D.E., Polin, D.S., Dunn, J.J., Wilske, B., Preac-Mursic, V., Dattwyler,
26 R.J., Luft, B.J., 1993. *Borrelia burgdorferi* is clonal: implications for taxonomy
27 and vaccine development. *Proc Natl Acad Sci U S A* 90, 10163-10167.
- 28 Eisen, L., Eisen, R.J., Lane, R.S., 2004. The roles of birds, lizards, and rodents as
29 hosts for the western black-legged tick *Ixodes pacificus*. *J Vector Ecol* 29, 295-
30 308.
- 31 Eisen, L., Eisen, R.J., Lane, R.S., 2006. Geographical distribution patterns and habitat
32 suitability models for presence of host-seeking ixodid ticks in dense woodlands of
33 Mendocino County, California. *J Med Entomol* 43, 415-427.
- 34 Eisen, L., Eisen, R.J., Mun, J., Salkeld, D.J., Lane, R.S., 2009. Transmission cycles of
35 *Borrelia burgdorferi* and *B. bissettii* in relation to habitat type in northwestern
36 California. *J Vector Ecol* 34, 81-91.
- 37 Eisen, L., Lane, R.S., 2002. Vectors of *Borrelia burgdorferi* sensu lato, in: Gray, J.,
38 Kahl, O., Lane, R.S., Stanek, G. (Eds.), *Lyme Borreliosis: Biology, Epidemiology*
39 *and Control*. CABI Publishing, Wallingford, pp. 91-115.
- 40 Excoffier, L., Heckel, G., 2006. Computer programs for population genetics data
41 analysis: a survival guide. *Nat Rev Genet* 7, 745-758.
- 42 Falco, R.C., Daniels, T.J., Fish, D., 1995. Increase in abundance of immature *Ixodes*
43 *scapularis* (Acari: Ixodidae) in an emergent Lyme disease endemic area. *J Med*
44 *Entomol* 32, 522-526.
- 45 Falco, R.C., Fish, D., 1991. Horizontal movement of adult *Ixodes dammini* (Acari:
46 Ixodidae) attracted to CO₂-baited traps. *J Med Entomol* 28, 726-729.
- 47 Feil, E.J., Cooper, J.E., Grundmann, H., Robinson, D.A., Enright, M.C., Berendt, T.,
48 Peacock, S.J., Smith, J.M., Murphy, M., Spratt, B.G., Moore, C.E., Day, N.P.,
49 2003. How clonal is *Staphylococcus aureus*? *J Bacteriol* 185, 3307-3316.

- 1 Feil, E.J., Enright, M.C., Spratt, B.G., 2000. Estimating the relative contributions of
2 mutation and recombination to clonal diversification: a comparison between
3 *Neisseria meningitidis* and *Streptococcus pneumoniae*. Res Microbiol 151, 465-
4 469.
- 5 Feil, E.J., Li, B.C., Aanensen, D.M., Hanage, W.P., Spratt, B.G., 2004. eBURST:
6 inferring patterns of evolutionary descent among clusters of related bacterial
7 genotypes from multilocus sequence typing data. J Bacteriol 186, 1518-1530.
- 8 Feil, E.J., Spratt, B.G., 2001. Recombination and the population structures of bacterial
9 pathogens. Annu Rev Microbiol 55, 561-590.
- 10 Fingerle, V., Michel, H., Hettche, G., Hizo-Teufel, C., Wilske, B., 2004. *Borrelia*
11 *burgdorferi* s.l. OspA-types are widespread in Bavaria but show distinct local
12 patterns. Int J Med Microbiol 293 Suppl 37, 165-166.
- 13 Francisco, A.P., Bugalho, M., Ramirez, M., Carrico, J.A., 2009. Global optimal
14 eBURST analysis of multilocus typing data using a graphic matroid approach.
15 BMC Bioinformatics 10, 152.
- 16 Frank, S.A., 2002. Immunology and Evolution of Infectious Disease. Princeton
17 University Press, Princeton.
- 18 Fukunaga, M., Hamase, A., 1995. Outer surface protein C gene sequence analysis of
19 *Borrelia burgdorferi* sensu lato isolates from Japan. J Clin Microbiol 33, 2415-
20 2420.
- 21 Fukunaga, M., Hamase, A., Okada, K., Inoue, H., Tsuruta, Y., Miyamoto, K., Nakao,
22 M., 1996a. Characterization of spirochetes isolated from ticks (*Ixodes tanuki*,
23 *Ixodes turdus*, and *Ixodes columnae*) and comparison of the sequences with those
24 of *Borrelia burgdorferi* sensu lato strains. Appl Environ Microbiol 62, 2338-2344.
- 25 Fukunaga, M., Hamase, A., Okada, K., Nakao, M., 1996b. *Borrelia tanukii* sp. nov.
26 and *Borrelia turdae* sp. nov. found from ixodid ticks in Japan: rapid species
27 identification by 16S rRNA gene-targeted PCR analysis. Microbiol Immunol 40,
28 877-881.
- 29 Fukunaga, M., Okada, K., Nakao, M., Konishi, T., Sato, Y., 1996c. Phylogenetic
30 analysis of *Borrelia* species based on flagellin gene sequences and its application
31 for molecular typing of Lyme disease borreliae. Int J Syst Bacteriol 46, 898-905.
- 32 Fukunaga, M., Takahashi, Y., Tsuruta, Y., Matsushita, O., Ralph, D., McClelland, M.,
33 Nakao, M., 1995. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov.,
34 isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in
35 Japan. Int J Syst Bacteriol 45, 804-810.
- 36 Gatewood, A.G., Liebman, K.A., Vourc'h, G., Bunikis, J., Hamer, S.A., Cortinas,
37 M.R., Melton, F., Cislo, P., Kitron, U., Tsao, J., Barbour, A.G., Fish, D., Diuk-
38 Wasser, M.A., 2009. Climate and tick seasonality predict *Borrelia burgdorferi*
39 genotype distribution. Applied and Environmental Microbiology 75, 2476-2483.
- 40 Gazumyan, A., Schwartz, J.J., Liveris, D., Schwartz, I., 1994. Sequence analysis of
41 the ribosomal RNA operon of the Lyme disease spirochete, *Borrelia burgdorferi*.
42 Gene 146, 57-65.
- 43 Gerlach, G., Musolf, K., 2000. Fragmentation of landscape as a cause for genetic
44 subdivision in bank voles. Conservation Biology 14, 1066-1074.
- 45 Gern, L., 2008. *Borrelia burgdorferi* sensu lato, the agent of lyme borreliosis: life in
46 the wilds. Parasite 15, 244-247.
- 47 Gern, L., Humair, P., 2002. Ecology of *Borrelia burgdorferi* sensu lato in Europe, in:
48 Gray, J.S., Kahl, O., Lane, R.S., Stanek, G. (Eds.), Lyme Borreliosis: Biology of
49 the Infectious Agents and Epidemiology of Disease CABI Publishing,
50 Wallingford, pp. 149-174.

1 Gern, L., Sell, K., 2009. Isolation of *Borrelia burgdorferi* sensu lato from the skin of
2 the European badger (*Meles meles*) in Switzerland. Vector Borne Zoonotic Dis 9,
3 207-208.

4 Gevers, D., Cohan, F.M., Lawrence, J.G., Spratt, B.G., Coenye, T., Feil, E.J.,
5 Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F.L., Swings, J.,
6 2005. Opinion: Re-evaluating prokaryotic species. Nat Rev Microbiol 3, 733-739.

7 Gevers, D., Dawyndt, P., Vandamme, P., Willems, A., Vancanneyt, M., Swings, J.,
8 De Vos, P., 2006. Stepping stones towards a new prokaryotic taxonomy. Philos
9 Trans R Soc Lond B Biol Sci 361, 1911-1916.

10 Girard, Y.A., Fedorova, N., Lane, R.S., 2011. Genetic diversity of *Borrelia*
11 *burgdorferi* and detection of *B. bissettii*-like DNA in serum of north-coastal
12 California residents. J Clin Microbiol 49, 945-954.

13 Girard, Y.A., Travinsky, B., Schotthoefer, A., Fedorova, N., Eisen, R.J., Eisen, L.,
14 Barbour, A.G., Lane, R.S., 2009. Population structure of the lyme borreliosis
15 spirochete *Borrelia burgdorferi* in the western black-legged tick (*Ixodes pacificus*)
16 in Northern California. Appl Environ Microbiol 75, 7243-7252.

17 Gomez-Diaz, E., Bouludier, T., Sertour, N., Cornet, M., Ferquel, E., McCoy, K.D.,
18 2011. Genetic structure of marine *Borrelia garinii* and population admixture with
19 the terrestrial cycle of Lyme borreliosis. Environ Microbiol.

20 Guillot, G., 2008. Inference of structure in subdivided populations at low levels of
21 genetic differentiation--the correlated allele frequencies model revisited.
22 Bioinformatics 24, 2222-2228.

23 Guillot, G., Santos, F., Estoup, A., 2008. Analysing georeferenced population genetics
24 data with Geneland: a new algorithm to deal with null alleles and a friendly
25 graphical user interface. Bioinformatics 24, 1406-1407.

26 Guttman, D.S., Stavrinides, J., 2010. Population Genomics of Bacteria, in: Robinson,
27 D.A., Falush, D., Feil, E.J. (Eds.), Bacterial Population Genetics in Infectious
28 Disease. John Wiley & Sons, Inc.

29 Hall, N., 2007. Advanced sequencing technologies and their wider impact in
30 microbiology. J Exp Biol 210, 1518-1525.

31 Hamer, S.A., Roy, P.L., Hickling, G.J., Walker, E.D., Foster, E.S., Barber, C.C.,
32 Tsao, J.I., 2007. Zoonotic pathogens in *Ixodes scapularis*, Michigan. Emerg Infect
33 Dis 13, 1131-1133.

34 Hamer, S.A., Tsao, J.I., Walker, E.D., Hickling, G.J., 2010. Invasion of the Lyme
35 Disease Vector *Ixodes scapularis*: Implications for *Borrelia burgdorferi*
36 Endemicity. Ecohealth DOI: 10.1007/s10393-010-0287-0.

37 Hanage, W.P., Fraser, C., Spratt, B.G., 2006. Sequences, sequence clusters and
38 bacterial species. Philos Trans R Soc Lond B Biol Sci 361, 1917-1927.

39 Hanage, W.P., Kaijalainen, T., Herva, E., Saukkoriipi, A., Syrjanen, R., Spratt, B.G.,
40 2005. Using multilocus sequence data to define the pneumococcus. J Bacteriol
41 187, 6223-6230.

42 Hanincova, K., Kurtenbach, K., Diuk-Wasser, M., Brei, B., Fish, D., 2006. Epidemic
43 spread of Lyme borreliosis, northeastern United States. Emerg Infect Dis 12, 604-
44 611.

45 Hanincova, K., Liveris, D., Sandigursky, S., Wormser, G.P., Schwartz, I., 2008a.
46 *Borrelia burgdorferi* sensu stricto is clonal in patients with early Lyme borreliosis.
47 Appl Environ Microbiol 74, 5008-5014.

48 Hanincova, K., Ogden, N.H., Diuk-Wasser, M., Pappas, C.J., Iyer, R., Fish, D.,
49 Schwartz, I., Kurtenbach, K., 2008b. Fitness variation of *Borrelia burgdorferi*
50 sensu stricto strains in mice. Appl Environ Microbiol 74, 153-157.

- 1 Hanincova, K., Schafer, S.M., Etti, S., Sewell, H.S., Taragelova, V., Ziak, D., Labuda,
2 M., Kurtenbach, K., 2003a. Association of *Borrelia afzelii* with rodents in Europe.
3 Parasitology 126, 11-20.
- 4 Hanincova, K., Taragelova, V., Koci, J., Schafer, S.M., Hails, R., Ullmann, A.J.,
5 Piesman, J., Labuda, M., Kurtenbach, K., 2003b. Association of *Borrelia garinii*
6 and *B. valaisiana* with songbirds in Slovakia. Appl Environ Microbiol 69, 2825-
7 2830.
- 8 Harris, S.R., Feil, E.J., Holden, M.T.G., Quail, M.A., Nickerson, E.K., Chanttrata,
9 N., Gardete, S., Tavares, A., Day, N., Lindsay, J.A., Edgeworth, J.D., Lencastre,
10 d.H., Parkhill, J., Paacock, S.J., Bentley, S.D., 2010. Evolution of MRSA During
11 Hospital Transmission and Intercontinental Spread. Science 327, 469-474.
- 12 Heckel, G., Burri, R., Fink, S., Desmet, J.F., Excoffier, L., 2005. Genetic structure
13 and colonization processes in European populations of the common vole,
14 *Microtus arvalis*. Evolution 59, 2231-2242.
- 15 Hellgren, O., Andersson, M., Raberg, L., 2011. The genetic structure of *Borrelia*
16 *afzelii* varies with geographic but not ecological sampling scale. J Evol Biol 24,
17 159-167.
- 18 Heroldova, M., Nemec, M., Hubalek, Z., 1998. Growth parameters of *Borrelia*
19 *burgdorferi* sensu stricto at various temperatures. Zentralbl Bakteriell 288, 451-
20 455.
- 21 Herrmann, C., Gern, L., 2010. Survival of *Ixodes ricinus* (Acari: Ixodidae) under
22 challenging conditions of temperature and humidity is influenced by *Borrelia*
23 *burgdorferi* sensu lato infection. J Med Entomol 47, 1196-1204.
- 24 Hewitt, G.M., 1999. Post-glacial re-colonization of European biota. Biological
25 Journal of the Linnean Society 68, 87-112.
- 26 Hewitt, G.M., 2001. Speciation, hybrid zones and phylogeography - or seeing genes
27 in space and time. Mol Ecol 10, 537-549.
- 28 Hoen, A.G., Margos, G., Bent, S.J., Kurtenbach, K., Fish, D., 2009. Phylogeography
29 of *Borrelia burgdorferi* in the eastern United States reveals multiple independent
30 Lyme disease emergence events. Proc Natl Acad Sci U S A 106, 15013-15018.
- 31 Holt, K.E., Parkhill, J., Mazzoni, C.J., Roumagnac, P., Weill, F.X., Goodhead, I.,
32 Rance, R., Baker, S., Maskell, D.J., Wain, J., Dolecek, C., Achtman, M., Dougan,
33 G., 2008. High-throughput sequencing provides insights into genome variation
34 and evolution in *Salmonella Typhi*. Nat Genet 40, 987-993.
- 35 Hu, C.M., Humair, P.F., Wallich, R., Gern, L., 1997. *Apodemus* sp. rodents, reservoir
36 hosts for *Borrelia afzelii* in an endemic area in Switzerland. Zentralbl Bakteriell
37 285, 558-564.
- 38 Hu, C.M., Wilske, B., Fingerle, V., Lobet, Y., Gern, L., 2001. Transmission of
39 *Borrelia garinii* OspA serotype 4 to BALB/c mice by *Ixodes ricinus* ticks
40 collected in the field. J Clin Microbiol 39, 1169-1171.
- 41 Hubalek, Z., Halouzka, J., 1997. Distribution of *Borrelia burgdorferi* sensu lato
42 genomic groups in Europe, a review. Eur J Epidemiol 13, 951-957.
- 43 Huegli, D., Hu, C.M., Humair, P.F., Wilske, B., Gern, L., 2002. *Apodemus* species
44 mice are reservoir hosts of *Borrelia garinii* OspA serotype 4 in Switzerland. J Clin
45 Microbiol 40, 4735-4737.
- 46 Huelsenbeck, J.P., Bull, J.J., Cunningham, C.W., 1996. Combining data in
47 phylogenetic analysis. Tree 11, 152-158.
- 48 Hulinska, D., Votypka, J., Kriz, B., Holinkova, N., Novakova, J., Hulinsky, V., 2007.
49 Phenotypic and genotypic analysis of *Borrelia* spp. isolated from *Ixodes ricinus*

ticks by using electrophoretic chips and real-time polymerase chain reaction. Folia Microbiol (Praha) 52, 315-324.

Humair, P., Gern, L., 2000. The wild hidden face of Lyme borreliosis in Europe. Microbes Infect 2, 915-922.

Humair, P.F., Postic, D., Wallich, R., Gern, L., 1998. An avian reservoir (*Turdus merula*) of the Lyme borreliosis spirochetes. Zentralbl Bakteriol 287, 521-538.

Humair, P.F., Rais, O., Gern, L., 1999. Transmission of *Borrelia afzelii* from *Apodemus* mice and *Clethrionomys* voles to *Ixodes ricinus* ticks: differential transmission pattern and overwintering maintenance. Parasitology 118 (Pt 1), 33-42.

Humphrey, P.T., Caporale, D.A., Brisson, D., 2010. Uncoordinated Phylogeography of *Borrelia burgdorferi* and Its Tick Vector, *Ixodes scapularis*. Evolution 64, 2653-2663.

Jauris-Heipke, S., Liegl, G., Preac-Mursic, V., Rossler, D., Schwab, E., Soutschek, E., Will, G., Wilske, B., 1995. Molecular analysis of genes encoding outer surface protein C (OspC) of *Borrelia burgdorferi* sensu lato: relationship to *ospA* genotype and evidence of lateral gene exchange of *ospC*. J Clin Microbiol 33, 1860-1866.

Johnson, R.C., Schmidt, G.P., Hyde, F.W., Steigerwalt, A.G., Brenner, D.J., 1984. *Borrelia burgdorferi* sp. nov.: etiological agent of Lyme disease. International Journal of Systematic Bacteriology 34, 496-497.

Kawabata, H., Masuzawa, T., Yanagihara, Y., 1993. Genomic analysis of *Borrelia japonica* sp. nov. isolated from *Ixodes ovatus* in Japan. Microbiol Immunol 37, 843-848.

Killilea, M.E., Swei, A., Lane, R.S., Briggs, C.J., Ostfeld, R.S., 2008. Spatial dynamics of lyme disease: a review. Ecohealth 5, 167-195.

Kollars, T.M., Jr., Oliver, J.H., Jr., Kollars, P.G., Durden, L.A., 1999. Seasonal activity and host associations of *Ixodes scapularis* (Acari: Ixodidae) in southeastern Missouri. J Med Entomol 36, 720-726.

Korenberg, E.I., Gorelova, N.B., Kovalevskii, Y.V., 2002. Ecology of *Borrelia burgdorferi* sensu lato in Russia, in: Gray, J., Kahl, O., Lane, R.S., Stanek, G. (Eds.), Lyme borreliosis: Biology, Epidemiology and Control. CABI Publishing, Wallingford.

Kuhner, M.K., 2006. LAMARC 2.0: maximum likelihood and Bayesian estimation of population parameters. Bioinformatics 22, 768-770.

Kurtenbach, K., De Michelis, S., Etti, S., Schafer, S.M., Sewell, H.S., Brade, V., Kraiczy, P., 2002b. Host association of *Borrelia burgdorferi* sensu lato-the key role of host complement. Trends Microbiol 10, 74-79.

Kurtenbach, K., De Michelis, S., Sewell, H.S., Etti, S., Schafer, S.M., Hails, R., Collares-Pereira, M., Santos-Reis, M., Hanincova, K., Labuda, M., Bormane, A., Donaghy, M., 2001. Distinct combinations of *Borrelia burgdorferi* sensu lato genospecies found in individual questing ticks from Europe. Appl Environ Microbiol 67, 4926-4929.

Kurtenbach, K., De Michelis, S., Sewell, H.S., Etti, S., Schafer, S.M., Holmes, E., Hails, R., Collares-Pereira, M., Santos-Reis, M., Hanincova, K., Labuda, M., Bormane, A., Donaghy, M., 2002. The key roles of selection and migration in the ecology of Lyme borreliosis. Int J Med Microbiol 291 Suppl 33, 152-154.

Kurtenbach, K., Hanincova, K., Tsao, J.I., Margos, G., Fish, D., Ogden, N.H., 2006. Fundamental processes in the evolutionary ecology of Lyme borreliosis. Nat Rev Microbiol 4, 660-669.

- 1 Kurtenbach, K., Hoen, A.G., Bent, S.J., Vollmer, S.A., Ogden, N.H., Margos, G.,
2 2010. Population Biology of Lyme Borreliosis spirochetes, in: Robinson, D.A.,
3 Falush, D., Feil, E.J. (Eds.), Bacterial Population Genetics in Infectious Disease, 1
4 ed. John Wiley & Sons, Inc.
- 5 Kurtenbach, K., Peacey, M., Rijpkema, S.G., Hoodless, A.N., Nuttall, P.A.,
6 Randolph, S.E., 1998a. Differential transmission of the genospecies of *Borrelia*
7 *burgdorferi* sensu lato by game birds and small rodents in England. Appl Environ
8 Microbiol 64, 1169-1174.
- 9 Kurtenbach, K., Schaefer, S.M., de Michelis, S., Etti, S., Sewell, H.S., 2002a.
10 *Borrelia burgdorferi* s.l. in the vertebrate host., in: Gray, J.S., Kahl, O., Lane,
11 R.S., Stanek, G. (Eds.), Lyme Borreliosis: Biology of the Infectious Agents and
12 Epidemiology of Disease. CABI Publishing, Wallingford, pp. 117-148.
- 13 Kurtenbach, K., Sewell, H.S., Ogden, N.H., Randolph, S.E., Nuttall, P.A., 1998b.
14 Serum complement sensitivity as a key factor in Lyme disease ecology. Infect
15 Immun 66, 1248-1251.
- 16 Lane, R.S., Quistad, G.B., 1998. Borreliacidal factor in the blood of the western fence
17 lizard (*Sceloporus occidentalis*). J Parasitol 84, 29-34.
- 18 Larsson, C., Comstedt, P., Olsen, B., Bergstrom, S., 2007. First record of Lyme
19 disease *Borrelia* in the Arctic. Vector Borne Zoonotic Dis 7, 453-456.
- 20 Le Fleche, A., Postic, D., Girardet, K., Peter, O., Baranton, G., 1997. Characterization
21 of *Borrelia lusitaniae* sp. nov. by 16S ribosomal DNA sequence analysis. Int J
22 Syst Bacteriol 47, 921-925.
- 23 Lin, T., Oliver, J.H., Jr., Gao, L., 2002. Genetic Diversity of the Outer Surface Protein
24 C Gene of Southern *Borrelia* Isolates and Its Possible Epidemiological, Clinical,
25 and Pathogenetic Implications. J Clin Microbiol 40, 2572-2583.
- 26 Lin, T., Oliver, J.H., Jr., Gao, L., 2003. Comparative analysis of *Borrelia* isolates
27 from southeastern USA based on randomly amplified polymorphic DNA
28 fingerprint and 16S ribosomal gene sequence analyses. FEMS Microbiol Lett 228,
29 249-257.
- 30 Liveris, D., Gazumyan, A., Schwartz, I., 1995. Molecular typing of *Borrelia*
31 *burgdorferi* sensu lato by PCR-restriction fragment length polymorphism analysis.
32 J Clin Microbiol 33, 589-595.
- 33 Loye, J.E., Lane, R.S., 1988. Questing behavior of *Ixodes pacificus* (Acari:Ixodidae)
34 in relation to meteorological and seasonal factors. Journal of Medical Entomology
35 25, 391-398.
- 36 Maddison, W.P., Knowles, L.L., 2006. Inferring phylogeny despite incomplete
37 lineage sorting. Syst Biol 55, 21-30.
- 38 Maggi, R.G., Reichelt, S., Toliver, M., Engber, B., 2010. *Borrelia* species in *Ixodes*
39 *affinis* and *Ixodes scapularis* ticks collected from the coastal plain of North
40 Carolina. Ticks Tick Borne Dis in press.
- 41 Maiden, M.C., 2006. Multilocus sequence typing of bacteria. Annu Rev Microbiol 60,
42 561-588.
- 43 Maiden, M.C., Bygraves, J.A., Feil, E., Morelli, G., Russell, J.E., Urwin, R., Zhang,
44 Q., Zhou, J., Zurth, K., Caugant, D.A., Feavers, I.M., Achtman, M., Spratt, B.G.,
45 1998. Multilocus sequence typing: a portable approach to the identification of
46 clones within populations of pathogenic microorganisms. Proc Natl Acad Sci U S
47 A 95, 3140-3145.
- 48 Manel, S., Schwartz, M.K., Luikart, G., Raberlet, P., 2003. Landscape genetics:
49 combining landscape ecology and populations genetics. Trends Ecology and
50 Evolution 18, 189-197.

- 1 Marconi, R.T., Garon, C.F., 1992. Phylogenetic analysis of the genus *Borrelia*: a
2 comparison of North American and European isolates of *Borrelia burgdorferi*. J
3 Bacteriol 174, 241-244.
- 4 Marconi, R.T., Liveris, D., Schwartz, I., 1995. Identification of novel insertion
5 elements, restriction fragment length polymorphism patterns, and discontinuous
6 23S rRNA in Lyme disease spirochetes: phylogenetic analyses of rRNA genes and
7 their intergenic spacers in *Borrelia japonica* sp. nov. and genomic group 21038
8 (*Borrelia andersonii* sp. nov.) isolates. J Clin Microbiol 33, 2427-2434.
- 9 Margos, G., Gatewood, A.G., Aanensen, D.M., Hanincova, K., Terekhova, D.,
10 Vollmer, S.A., Cornet, M., Piesman, J., Donaghy, M., Bormane, A., Hurn, M.A.,
11 Feil, E.J., Fish, D., Casjens, S., Wormser, G.P., Schwartz, I., Kurtenbach, K.,
12 2008. MLST of housekeeping genes captures geographic population structure and
13 suggests a European origin of *Borrelia burgdorferi*. Proc Natl Acad Sci U S A
14 105, 8730-8735.
- 15 Margos, G., Hojgaard, A., Lane, R.S., Cornet, M., Fingerle, V., Rudenko, N., Ogden,
16 N., Aanensen, D.M., Fish, D., Piesman, J., 2010. Multilocus sequence analysis of
17 *Borrelia bissettii* strains from North America reveals a new *Borrelia* species,
18 *Borrelia kurtenbachii*. Ticks Tick Borne Dis 1, 151-158.
- 19 Margos, G., Vollmer, S.A., Cornet, M., Garnier, M., Fingerle, V., Wilske, B.,
20 Bormane, A., Vitorino, L., Collares-Pereira, M., Drancourt, M., Kurtenbach, K.,
21 2009. A new *Borrelia* species defined by Multilocus Sequence Analysis of
22 Housekeeping Genes. Appl Environ Microbiol 75, 5410-5416.
- 23 Marti Ras, N., Postic, D., Foretz, M., Baranton, G., 1997. *Borrelia burgdorferi* sensu
24 stricto, a bacterial species "made in the U.S.A."? Int J Syst Bacteriol 47, 1112-
25 1117.
- 26 Masuzawa, T., 2004. Terrestrial distribution of the Lyme borreliosis agent *Borrelia*
27 *burgdorferi* sensu lato in East Asia. Jpn J Infect Dis 57, 229-235.
- 28 Masuzawa, T., Kharitonov, I.G., Kadosaka, T., Hashimoto, N., Kudeken, M.,
29 Takada, N., Kaneda, K., Imai, Y., 2005. Characterization of *Borrelia burgdorferi*
30 sensu lato isolated in Moscow province-a sympatric region for *Ixodes ricinus* and
31 *Ixodes persulcatus*. Int J Med Microbiol 294, 455-464.
- 32 Mather, T.N., Fish, D., Coughlin, R.T., 1994. Competence of dogs as reservoirs for
33 Lyme disease spirochetes (*Borrelia burgdorferi*). J Am Vet Med Assoc 205, 186-
34 188.
- 35 Mathiesen, D.A., Oliver, J.H., Jr., Kolbert, C.P., Tullson, E.D., Johnson, B.J.,
36 Campbell, G.L., Mitchell, P.D., Reed, K.D., Telford, S.R., 3rd, Anderson, J.F.,
37 Lane, R.S., Persing, D.H., 1997. Genetic heterogeneity of *Borrelia burgdorferi* in
38 the United States. J Infect Dis 175, 98-107.
- 39 Matuschka, F.R., Schinkel, T.W., Klug, B., Spielman, A., Richter, D., 2000. Relative
40 incompetence of European rabbits for Lyme disease spirochaetes. Parasitology
41 121, 297-302.
- 42 Maupin, G.O., Gage, K.L., Piesman, J., Montenieri, J., Sviat, S.L., VanderZanden, L.,
43 Happ, C.M., Dolan, M., Johnson, B.J., 1994. Discovery of an enzootic cycle of
44 *Borrelia burgdorferi* in *Neotoma mexicana* and *Ixodes spinipalpis* from northern
45 Colorado, an area where Lyme disease is nonendemic. J Infect Dis 170, 636-643.
- 46 McCabe, T.R., McCabe, R.E., 1997. Recounting whitetails past, in: McShea, W.J.,
47 Underwood, H.B., Rappole, J.H. (Eds.), The Science of Overabundance: Deer
48 Ecology and Population Management. Smithsonian Institution Press, Washington
49 DC, pp. 11-26.

- 1 Michel, H., Wilske, B., Hettche, G., Gottner, G., Heimerl, C., Reischl, U., Schulte-
2 Spechtel, U., Fingerle, V., 2004. An *ospA*-polymerase chain reaction/restriction
3 fragment length polymorphism-based method for sensitive detection and reliable
4 differentiation of all European *Borrelia burgdorferi* sensu lato species and OspA
5 types. Med Microbiol Immunol 193, 219-226.
- 6 Miyamoto, K., Masuzawa, T., 2002. Ecology of *Borrelia burgdorferi* sensu lato in
7 Japan and East Asia, in: Gray, J., Kahl, O., Lane, R.S., Stanek, G. (Eds.), Lyme
8 Borreliosis: Biology, Epidemiology and Control. CABI Publishing, Wallingford,
9 pp. 201-222.
- 10 Murphy, K.M., Geiger, T., Hafez, M.J., Eshleman, J.R., Griffin, C.A., Berg, K.D.,
11 2003. A single nucleotide primer extension assay to detect the APC I1307K gene
12 variant. J Mol Diagn 5, 222-226.
- 13 Nefedova, V.V., Korenberg, E.I., Gorelova, N.B., Kovalevskii, Y.V., 2004. Studies
14 on the transovarial transmission of *Borrelia burgdorferi* sensu lato in the taiga tick
15 *Ixodes persulcatus* Folia Parasitology 51 67-71.
- 16 Norris, D.E., Johnson, B.J., Piesman, J., Maupin, G.O., Clark, J.L., Black, W.C.t.,
17 1999. Population genetics and phylogenetic analysis of Colorado *Borrelia*
18 *burgdorferi*. Am J Trop Med Hyg 60, 699-707.
- 19 Norris, D.E., Klompen, J.S., Keirans, J.E., Black, W.C.t., 1996. Population genetics of
20 *Ixodes scapularis* (Acari: Ixodidae) based on mitochondrial 16S and 12S genes. J
21 Med Entomol 33, 78-89.
- 22 Ochman, H., Lawrence, J.G., Gorisman, E.A., 2000. Lateral gene transfer and the
23 nature of bacterial innovation. Nature 405, 299-304.
- 24 Ogden, N.H., Bigras-Poulin, M., Hanincova, K., Maarouf, A., O'Callaghan, C.J.,
25 Kurtenbach, K., 2008a. Projected effects of climate change on tick phenology and
26 fitness of pathogens transmitted by the North American tick *Ixodes scapularis*. J
27 Theor Biol 254, 621-632.
- 28 Ogden, N.H., Bigras-Poulin, M., O'Callaghan C, J., Barker, I.K., Kurtenbach, K.,
29 Lindsay, L.R., Charron, D.F., 2007. Vector seasonality, host infection dynamics
30 and fitness of pathogens transmitted by the tick *Ixodes scapularis*. Parasitology
31 134, 209-227.
- 32 Ogden, N.H., Bouchard, C., Kurtenbach, K., Margos, G., Lindsay, L.R., Trudel, L.,
33 Nguon, S., Milord, F., 2010. Active and passive surveillance and phylogenetic
34 analysis of *Borrelia burgdorferi* elucidate the process of Lyme disease risk
35 emergence in Canada. Environ Health Perspect 118, 909-914.
- 36 Ogden, N.H., Lindsay, L.R., Hanincova, K., Barker, I.K., Bigras-Poulin, M., Charron,
37 D.F., Heagy, A., Francis, C.M., O'Callaghan, C.J., Schwartz, I., Thompson, R.A.,
38 2008b. Role of migratory birds in introduction and range expansion of *Ixodes*
39 *scapularis* ticks and of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in
40 Canada. Appl Environ Microbiol 74, 1780-1790.
- 41 Ogden, N.H., Lindsay, L.R., Morshed, M., Sockett, P.N., Artsob, H., 2009a. The
42 emergence of Lyme disease in Canada. CMAJ Canadian Medical Association
43 Journal 180, 1221-1224.
- 44 Ogden, N.H., Margos, G., Aanensen, D.M., Drebot, M.A., Feil, E.J., Hanincová, K.,
45 Schwartz, I., Tyler, S., Lindsay, L.R., 2011. Investigation of genotypes of *Borrelia*
46 *burgdorferi* in *Ixodes scapularis* ticks collected in surveillance in Canada. Appl
47 Environ Microbiol.
- 48 Ogden, N.H., Nuttall, P.A., Randolph, S.E., 1997. Natural Lyme disease cycles
49 maintained via sheep by co-feeding ticks. Parasitology 115 (Pt 6), 591-599.

- 1 Ogden, N.H., Trudel, L., Artsob, H., Barker, I.K., Beauchamp, G., Charron, D.F.,
2 Drebot, M.A., Galloway, T.D., O'Handley, R., Thompson, R.A., Lindsay, L.R.,
3 2006. *Ixodes scapularis* ticks collected by passive surveillance in Canada: analysis
4 of geographic distribution and infection with Lyme borreliosis agent *Borrelia*
5 *burgdorferi*. J Med Entomol 43, 600-609.
- 6 Ogden, N.H., Tsao, J.I., 2009b. Biodiversity and Lyme disease: dilution or
7 amplification? Epidemics 1, 196-206.
- 8 Oliver, J., Means, R.G., Kogut, S., Prusinski, M., Howard, J.J., Layne, L.J., Chu, F.K.,
9 Reddy, A., Lee, L., White, D.J., 2006. Prevalence of *Borrelia burgdorferi* in small
10 mammals in New York state. J Med Entomol 43, 924-935.
- 11 Oliver, J.H., Jr., 1996. Lyme borreliosis in the southern United States: a review. J
12 Parasitol 82, 926-935.
- 13 Oliver, J.H., Jr., Lin, T., Gao, L., Clark, K.L., Banks, C.W., Durden, L.A., James,
14 A.M., Chandler, F.W., Jr., 2003. An enzootic transmission cycle of Lyme
15 borreliosis spirochetes in the southeastern United States. Proc Natl Acad Sci U S
16 A 100, 11642-11645.
- 17 Olsen, B., Duffy, D.C., Jaenson, T.G., Gylfe, A., Bonnedahl, J., Bergstrom, S., 1995.
18 Transhemispheric exchange of Lyme disease spirochetes by seabirds. J Clin
19 Microbiol 33, 3270-3274.
- 20 Olsen, B., Jaenson, T.G., Noppa, L., Bunikis, J., Bergstrom, S., 1993. A Lyme
21 borreliosis cycle in seabirds and *Ixodes uriae* ticks. Nature 362, 340-342.
- 22 Ornstein, K., Berglund, J., Nilsson, I., Norrby, R., Bergstrom, S., 2001.
23 Characterization of Lyme borreliosis isolates from patients with erythema migrans
24 and neuroborreliosis in southern Sweden. J Clin Microbiol 39, 1294-1298.
- 25 Page, R.D.M., Holmes, E.C., 1998. Molecular Evolution: a phylogenetic approach.
26 Blackwell Publishing Ltd., Oxford.
- 27 Park, H.S., Lee, J.H., Jeong, E.J., Koh, S.E., Park, T.K., Jang, W.J., Park, K.H., Kim,
28 B.J., Kook, Y.H., Lee, S.H., 2004. Evaluation of *groEL* gene analysis for
29 identification of *Borrelia burgdorferi* sensu lato. J Clin Microbiol 42, 1270-1273.
- 30 Patrican, L.A., 1997. Absence of Lyme disease spirochetes in larval progeny of
31 naturally infected *Ixodes scapularis* (Acari: Ixodidae) fed on dogs. Journal of
32 Medical Entomology 34 52-55.
- 33 Paulo, O.S., Pinheiro, J., Miraldo, A., Bruford, M.W., Jordan, W.C., Nichols, R.A.,
34 2008. The role of vicariance vs. dispersal in shaping genetic patterns in ocellated
35 lizard species in the western Mediterranean. Molecular Ecology 17, 1535-1551.
- 36 Picken, R.N., Cheng, Y., Han, D., Nelson, J.A., Reddy, A.G., Hayden, M.K., Picken,
37 M.M., Strle, F., Bouseman, J.K., Trenholme, G.M., 1995. Genotypic and
38 phenotypic characterization of *Borrelia burgdorferi* isolated from ticks and small
39 animals in Illinois. J Clin Microbiol 33, 2304-2315.
- 40 Picken, R.N., Cheng, Y., Strle, F., Cimperman, J., Maraspin, V., Lotric-Furlan, S.,
41 Ruzic-Sabljic, E., Han, D., Nelson, J.A., Picken, M.M., Trenholme, G.M., 1996a.
42 Molecular characterization of *Borrelia burgdorferi* sensu lato from Slovenia
43 revealing significant differences between tick and human isolates. Eur J Clin
44 Microbiol Infect Dis 15, 313-323.
- 45 Picken, R.N., Cheng, Y., Strle, F., Picken, M.M., 1996b. Patient isolates of *Borrelia*
46 *burgdorferi* sensu lato with genotypic and phenotypic similarities of strain 25015.
47 J Infect Dis 174, 1112-1115.
- 48 Picken, R.N., Picken, M.M., 2000. Molecular characterization of *Borrelia* spp.
49 isolates from greater metropolitan Chicago reveals the presence of *Borrelia*
50 *bissettii*. Preliminary report. J Mol Microbiol Biotechnol 2, 505-507.

- 1 Piesman, J., 1993. Standard system for infecting ticks (Acari: Ixodidae) with the
2 Lyme disease spirochete, *Borrelia burgdorferi*. J Med Entomol 30, 199-203.
- 3 Piesman, J., 2002. Ecology of *Borrelia burgdorferi* sensu lato in Northamerica in:
4 Gray, J.S., Kahl, O., Lane, R.S., Stanek, G. (Eds.), Lyme Borreliosis: Biology of
5 the Infectious Agents and Epidemiology of Disease CABI Publishing,
6 Wallingford, pp. 223-249.
- 7 Piesman, J., Gern, L., 2004. Lyme borreliosis in Europe and North America.
8 Parasitology 129 Suppl, S191-220.
- 9 Piesman, J., Schwan, T.G., 2010. Ecology of borreliae and their arthropod vectors in:
10 Samuels, D.S., Radolf, J.D. (Eds.), *Borrelia: Molecular Biology, Host Interaction*
11 *and Pathogenesis*. Caister Academic Press, pp. 251-278.
- 12 Pollack, R.J., Telford, S.R., 3rd, Spielman, A., 1993. Standardization of medium for
13 culturing Lyme disease spirochetes. J Clin Microbiol 31, 1251-1255.
- 14 Portnoi, D., Sertour, N., Ferquel, E., Garnier, M., Baranton, G., Postic, D., 2006. A
15 single-run, real-time PCR for detection and identification of *Borrelia burgdorferi*
16 sensu lato species, based on the *hbb* gene sequence. FEMS Microbiol Lett 259,
17 35-40.
- 18 Postic, D., Assous, M.V., Grimont, P.A., Baranton, G., 1994. Diversity of *Borrelia*
19 *burgdorferi* sensu lato evidenced by restriction fragment length polymorphism of
20 rrf (5S)-rrl (23S) intergenic spacer amplicons. Int J Syst Bacteriol 44, 743-752.
- 21 Postic, D., Garnier, M., Baranton, G., 2007. Multilocus sequence analysis of atypical
22 *Borrelia burgdorferi* sensu lato isolates - description of *Borrelia californiensis* sp.
23 nov., and genomospecies 1 and 2. Int J Med Microbiol 297, 263-271.
- 24 Postic, D., Ras, N.M., Lane, R.S., Henderson, M., Baranton, G., 1998. Expanded
25 diversity among Californian borrelia isolates and description of *Borrelia bissettii*
26 sp. nov. (formerly *Borrelia* group DN127). J Clin Microbiol 36, 3497-3504.
- 27 Qiu, W.G., Bosler, E.M., Campbell, J.R., Ugine, G.D., Wang, I.N., Luft, B.J.,
28 Dykhuizen, D.E., 1997. A population genetic study of *Borrelia burgdorferi* sensu
29 stricto from eastern Long Island, New York, suggested frequency-dependent
30 selection, gene flow and host adaptation. Hereditas 127, 203-216.
- 31 Qiu, W.G., Bruno, J.F., McCaig, W.D., Xu, Y., Livey, I., Schriefer, M.E., Luft, B.J.,
32 2008. Wide distribution of a high-virulence *Borrelia burgdorferi* clone in Europe
33 and North America. Emerg Infect Dis 14, 1097-1104.
- 34 Qiu, W.G., Dykhuizen, D.E., Acosta, M.S., Luft, B.J., 2002. Geographic uniformity
35 of the Lyme disease spirochete (*Borrelia burgdorferi*) and its shared history with
36 tick vector (*Ixodes scapularis*) in the Northeastern United States. Genetics 160,
37 833-849.
- 38 Qiu, W.G., Schutzer, S.E., Bruno, J.F., Attie, O., Xu, Y., Dunn, J.J., Fraser, C.M.,
39 Casjens, S.R., Luft, B.J., 2004. Genetic exchange and plasmid transfers in
40 *Borrelia burgdorferi* sensu stricto revealed by three-way genome comparisons and
41 multilocus sequence typing. Proc Natl Acad Sci U S A 101, 14150-14155.
- 42 Randolph, S.E., 1998. Ticks are not Insects: Consequences of Contrasting Vector
43 Biology for Transmission Potential. Parasitol Today 14, 186-192.
- 44 Randolph, S.E., 2008. Dynamics of tick-borne disease systems: minor role of recent
45 climate change. Rev Sci Tech 27, 367-381.
- 46 Rauter, C., Hartung, T., 2005. Prevalence of *Borrelia burgdorferi* sensu lato
47 genospecies in *Ixodes ricinus* ticks in Europe: A metaanalysis. Appl Environ
48 Microbiol 71, 7203-7216.
- 49 Richter, D., Matuschka, F.R., 2006. Perpetuation of the Lyme disease spirochete
50 *Borrelia lusitaniae* by lizards. Appl Environ Microbiol 72, 4627-4632.

- 1 Richter, D., Postic, D., Sertour, N., Livey, I., Matuschka, F.R., Baranton, G., 2006.
- 2 Delineation of *Borrelia burgdorferi* sensu lato species by multilocus sequence
- 3 analysis and confirmation of the delineation of *Borrelia spielmanii* sp. nov. Int J
- 4 Syst Evol Microbiol 56, 873-881.
- 5 Richter, D., Spielman, A., Komar, N., Matuschka, F.R., 2000. Competence of
- 6 American robins as reservoir hosts for Lyme disease spirochetes. Emerg Infect
- 7 Dis 6, 133-138.
- 8 Rijpkema, S.G., Tazelaar, D.J., Molkenboer, M.J., Noordhoek, G.T., Plantinga, G.,
- 9 Schouls, L.M., Schellekens, J.F., 1997. Detection of *Borrelia afzelii*, *Borrelia*
- 10 *burgdorferi* sensu stricto, *Borrelia garinii* and group VS116 by PCR in skin
- 11 biopsies of patients with erythema migrans and acrodermatitis chronica
- 12 atrophicans. Clin Microbiol Infect 3, 109-116.
- 13 Rosa, P.A., Schwan, T., Hogan, D., 1992. Recombination between genes encoding
- 14 major outer surface proteins A and B of *Borrelia burgdorferi*. Molecular
- 15 Microbiology 6, 3031-3040.
- 16 Rudenko, N., Golovchenko, M., Grubhoffer, L., Oliver, J.H., Jr., 2009a. *Borrelia*
- 17 *carolinensis* sp.nov. - a new (14th) member of *Borrelia burgdorferi* sensu lato
- 18 complex from the southeastern United States. J Clin Microbiol 47, 134-141.
- 19 Rudenko, N., Golovchenko, M., Lin, T., Gao, L., Grubhoffer, L., Oliver, J.H., Jr.,
- 20 2009b. Delineation of a new species of the *Borrelia burgdorferi* sensu lato
- 21 complex, *Borrelia americana* sp.nov. J Clin Microbiol 47, 3875-3880.
- 22 Rudenko, N., Golovchenko, M., Mokracek, A., Piskunova, N., Ruzek, D., Mallatova,
- 23 N., Grubhoffer, L., 2008. Detection of *Borrelia bissettii* in cardiac valve tissue of
- 24 a patient with endocarditis and aortic valve stenosis in the Czech Republic. J Clin
- 25 Microbiol 46, 3540-3543.
- 26 Sadziene, A., Wilske, B., Ferdows, M.S., Barbour, A.G., 1993. The cryptic *ospC* gene
- 27 of *Borrelia burgdorferi* B31 is located on a circular plasmid. Infect Immun 61,
- 28 2192-2195.
- 29 Schierup, M.H., Wiuf, C., 2010. The coalescent of bacterial populations, in:
- 30 Robinson, D.A., Falush, D., Feil, E.J. (Eds.), Bacterial Population Genetics in
- 31 Infectious Disease. John Wiley & Sons, Inc., Hoboken, NJ.
- 32 Schulte-Spechtel, U., Fingerle, V., Goettner, G., Rogge, S., Wilske, B., 2006.
- 33 Molecular analysis of decorin-binding protein A (DbpA) reveals five major
- 34 groups among European *Borrelia burgdorferi* sensu lato strains with impact for
- 35 the development of serological assays and indicates lateral gene transfer of the
- 36 *dbpA* gene. Int J Med Microbiol 296 Suppl 40, 250-266.
- 37 Schutzer, S.E., Fraser-Liggett, C.M., Casjens, S.R., Qiu, W.G., Dunn, J.J., Mongodin,
- 38 E.F., Luft, B.J., 2011. Whole-genome sequences of thirteen isolates of *Borrelia*
- 39 *burgdorferi*. J Bacteriol 193, 1018-1020.
- 40 Schwartz, J.J., Gazumyan, A., Schwartz, I., 1992. rRNA gene organization in the
- 41 Lyme disease spirochete, *Borrelia burgdorferi*. J Bacteriol 174, 3757-3765.
- 42 Schweizer, M., Excoffier, L., Heckel, G., 2007. Fine-scale genetic structure and
- 43 dispersal in the common vole (*Microtus arvalis*). Mol Ecol 16, 2463-2473.
- 44 Scoles, G.A., Papero, M., Beati, L., Fish, D., 2001. A relapsing fever group spirochete
- 45 transmitted by *Ixodes scapularis* ticks. Vector Borne Zoonotic Dis 1, 21-34.
- 46 Scott, J.D., Lee, M.K., Fernando, K., Durden, L.A., Jorgensen, D.R., Mak, S.,
- 47 Morshed, M.G., 2010. Detection of Lyme disease spirochete, *Borrelia burgdorferi*
- 48 sensu lato, including three novel genotypes in ticks (Acari: Ixodidae) collected
- 49 from songbirds (Passeriformes) across Canada. J Vector Ecol 35, 124-139.

- 1 Searle, J.B., Kotlik, P., Rambau, R.V., Markova, S., Herman, J.S., McDevitt, A.D.,
2 2009. The Celtic fringe of Britain: insights from small mammal phylogeography.
3 Proc Biol Sci 276, 4287-4294.
- 4 Seinost, G., Dykhuizen, D.E., Dattwyler, R.J., Golde, W.T., Dunn, J.J., Wang, I.N.,
5 Wormser, G.P., Schriefer, M.E., Luft, B.J., 1999. Four clones of *Borrelia*
6 *burgdorferi* sensu stricto cause invasive infection in humans. Infect Immun 67,
7 3518-3524.
- 8 Smith, R.P., Jr., Muzaffar, S.B., Lavers, J., Lacombe, E.H., Cahill, B.K., Lubelczyk,
9 C.B., Kinsler, A., Mathers, A.J., Rand, P.W., 2006. *Borrelia garinii* in seabird
10 ticks (*Ixodes uriae*), Atlantic Coast, North America. Emerg Infect Dis 12, 1909-
11 1912.
- 12 Spielman, A., 1994. The emergence of Lyme disease and human babesiosis in a
13 changing environment. Ann N Y Acad Sci 740, 146-156.
- 14 Spielman, A., Levine, J.F., Wilson, M.L., 1984. Vectorial capacity of North American
15 *Ixodes* ticks. Yale J Biol Med 57, 507-513.
- 16 Stackebrandt, E., Ebers, J., 2006. Taxonomic parameters revisited: tarnished gold
17 standards. Microbiology Today 33, 152-155.
- 18 Staley, J.T., 2006. The bacterial species dilemma and the genomic-phylogenetic
19 species concept. Philos Trans R Soc Lond B Biol Sci 361, 1899-1909.
- 20 Stanek, G., Strle, F., 2009. Lyme borreliosis: a European perspective on diagnosis and
21 clinical management. Curr Opin Infect Dis 22, 450-454.
- 22 Steere, A.C., Bartenhagen, N.H., Craft, J.E., Hutchinson, G.J., Newman, J.H.,
23 Pachner, A.R., Rahn, D.W., Sigal, L.H., Taylor, E., Malawista, S.E., 1986.
24 Clinical manifestations of Lyme disease. Zentralbl Bakteriol Mikrobiol Hyg [A]
25 263, 201-205.
- 26 Strube, C., Montenegro, V.M., Epe, C., Eckelt, E., Schnieder, T., 2010. Establishment
27 of a minor groove binder-probe based quantitative real time PCR to detect
28 *Borrelia burgdorferi* sensu lato and differentiation of *Borrelia spielmanii* by
29 ospA-specific conventional PCR. Parasit Vectors 3, 69.
- 30 Swanson, K.I., Norris, D.E., 2007. Detection of *Borrelia burgdorferi* DNA in lizards
31 from Southern Maryland. Vector Borne Zoonotic Dis 7, 42-49.
- 32 Swei, A., Ostfeld, R.S., Lane, R.S., Briggs, C.J., 2011. Impact of the experimental
33 removal of lizards on Lyme disease risk. Proc Biol Sci.
- 34 Taberlet, P., Bouvet, J., 1994. Mitochondrial DNA polymorphism, phylogeography,
35 and conservation genetics of the brown bear *Ursus arctos* in Europe. Proc Biol Sci
36 255, 195-200.
- 37 Taberlet, P., Fumagalli, L., Wust-Saucy, A.G., Cosson, J.F., 1998. Comparative
38 phylogeography and postglacial colonization routes in Europe. Mol Ecol 7, 453-
39 464.
- 40 Takano, A., Nakao, M., Masuzawa, T., Takada, N., Yano, Y., Ishiguro, F., Fujita, H.,
41 Ito, T., Ma, X., Oikawa, Y., Kawamori, F., Kumagai, K., Mikami, T., Hanaoka,
42 N., Ando, S., Honda, N., Taylor, K., Tsubota, T., Konnai, S., Watanabe, H.,
43 Ohnishi, M., Kawabata, H., 2011. Multilocus Sequence Typing Implicates
44 Rodents as the Main Reservoir Host of Human-Pathogenic *Borrelia garinii* in
45 Japan. J Clin Microbiol 49, 2035-2039.
- 46 Talleklint-Eisen, L., Eisen, R.J., 1999. Abundance of ticks (Acari: Ixodidae) infesting
47 the western fence lizard, *Sceloporus occidentalis*, in relation to environmental
48 factors. Exp Appl Acarol 23, 731-740.
- 49 Taragel'ova, V., Koci, J., Hanincova, K., Kurtenbach, K., Derdakova, M., Ogden,
50 N.H., Literak, I., Kocianova, E., Labuda, M., 2008. Blackbirds and song thrushes

constitute a key reservoir of *Borrelia garinii*, the causative agent of borreliosis in Central Europe. Appl Environ Microbiol 74, 1289-1293.

Telford, S.R., 3rd, Mather, T.N., Moore, S.I., Wilson, M.L., Spielman, A., 1988. Incompetence of deer as reservoirs of the Lyme disease spirochete. Am J Trop Med Hyg 39, 105-109.

Theisen, M., Frederiksen, B., Lebech, A.M., Vuust, J., Hansen, K., 1993. Polymorphism in *ospC* gene of *Borrelia burgdorferi* and immunoreactivity of OspC protein: implications for taxonomy and for use of OspC protein as a diagnostic antigen. J Clin Microbiol 31, 2570-2576.

Travinsky, B., Bunikis, J., Barbour, A.G., 2010. Geographic differences in genetic locus linkages for *Borrelia burgdorferi*. Emerg Infect Dis 16, 1147-1150.

Tsao, J.I., 2009. Reviewing molecular adaptations of Lyme borreliosis spirochetes in the context of reproductive fitness in natural transmission cycles. Vet Res 40, 36.

Ullmann, A.J., Lane, R.S., Kurtenbach, K., Miller, M., Schriefer, M.E., Zeldner, N., Piesman, J., 2003. Bacteriolytic activity of selected vertebrate sera for *Borrelia burgdorferi* sensu stricto and *Borrelia bissettii*. J Parasitol 89, 1256-1257.

Urwin, R., Maiden, M.C., 2003. Multi-locus sequence typing: a tool for global epidemiology. Trends Microbiol 11, 479-487.

Valsangiacomo, C., Balmelli, T., Piffaretti, J.C., 1997. A phylogenetic analysis of *Borrelia burgdorferi* sensu lato based on sequence information from the *hbb* gene, coding for a histone-like protein. Int J Syst Bacteriol 47, 1-10.

van Dam, A.P., 2002. Diversity of *Ixodes*-borne *Borrelia* species - clinical, pathogenetic, and diagnostic implications and impact on vaccine development. Vector Borne Zoonotic Dis 2, 249-254.

Vitorino, L.R., Margos, G., Feil, E.J., Collares-Pereira, M., Ze-Ze, L., Kurtenbach, K., 2008. Fine-scale Phylogeographic Structure of *Borrelia lusitaniae* Revealed by Multilocus Sequence Typing. PloS ONE 3, e4002.

Vollmer, S.A., Margos, G., Donaghy, M., Bormane, A., Drancourt, M., Garnier, M., Cornet, M., Kurtenbach, K., 2010. Phylogeographic Structuring and Evolutionary Relationships of Lyme Borreliosis Spirochetes in Europe as Revealed by MLSA. Environmental Microbiology 13, 184-192.

Vollmer, S.A., Margos, G., Donaghy, M., Bormane, A., Drancourt, M., Garnier, M., Cornet, M., Kurtenbach, K., 2011. Phylogeographic Structuring and Evolutionary Relationships of Lyme Borreliosis Spirochetes in Europe as Revealed by MLSA. Environmental Microbiology 13, 184-192.

Wang, G., van Dam, A.P., Dankert, J., 1999a. Evidence for frequent OspC gene transfer between *Borrelia valaisiana* sp. nov. and other Lyme disease spirochetes. FEMS Microbiol Lett 177, 289-296.

Wang, G., van Dam, A.P., Dankert, J., 2000. Two distinct *ospA* genes among *Borrelia valaisiana* strains. Res Microbiol 151, 325-331.

Wang, G., van Dam, A.P., Le Fleche, A., Postic, D., Peter, O., Baranton, G., de Boer, R., Spanjaard, L., Dankert, J., 1997. Genetic and phenotypic analysis of *Borrelia valaisiana* sp. nov. (*Borrelia* genomic groups VS116 and M19). Int J Syst Bacteriol 47, 926-932.

Wang, G., van Dam, A.P., Schwartz, I., Dankert, J., 1999b. Molecular typing of *Borrelia burgdorferi* sensu lato: taxonomic, epidemiological, and clinical implications. Clin Microbiol Rev 12, 633-653.

Wang, I.N., Dykhuizen, D.E., Qiu, W., Dunn, J.J., Bosler, E.M., Luft, B.J., 1999. Genetic diversity of *ospC* in a local population of *Borrelia burgdorferi* sensu stricto. Genetics 151, 15-30.

- 1 Wayne, L.G., Brenner, D.J., Colwell, R.R., Grimont, R.A.D., Moore, W.E.C.,
2 Murray, R.G.E., Stackebrandt, E., Starrm, P., Truper, H.G., 1987. Report of the ad
3 hoe committee on reconciliation of approaches to bacterial systematics.
4 International Journal of Systematic Bacteriology 37, 463-464.
- 5 Will, G., Jauris-Heipke, S., Schwab, E., Busch, U., Rossler, D., Soutschek, E.,
6 Wilske, B., Preac-Mursic, V., 1995. Sequence analysis of *ospA* genes shows
7 homogeneity within *Borrelia burgdorferi* sensu stricto and *Borrelia afzelii* strains
8 but reveals major subgroups within the *Borrelia garinii* species. Med Microbiol
9 Immunol 184, 73-80.
- 10 Wilske, B., Anderson, J.F., Baranton, G., Barbour, A.G., Hovind-Hougen, K.,
11 Johnson, R.C., Preac-Mursic, V., 1991. Taxonomy of *Borrelia* spp. Scand J Infect
12 Dis Suppl 77, 108-129.
- 13 Wilske, B., Busch, U., Eiffert, H., Fingerle, V., Pfister, H.W., Rossler, D., Preac-
14 Mursic, V., 1996. Diversity of OspA and OspC among cerebrospinal fluid isolates
15 of *Borrelia burgdorferi* sensu lato from patients with neuroborreliosis in
16 Germany. Med Microbiol Immunol 184, 195-201.
- 17 Wilske, B., Jauris-Heipke, S., Lobentanzer, R., Pradel, I., Preac-Mursic, V., Rossler,
18 D., Soutschek, E., Johnson, R.C., 1995. Phenotypic analysis of outer surface
19 protein C (OspC) of *Borrelia burgdorferi* sensu lato by monoclonal antibodies:
20 relationship to genospecies and OspA serotype. J Clin Microbiol 33, 103-109.
- 21 Wilske, B., Preac-Mursic, V., Gobel, U.B., Graf, B., Jauris, S., Soutschek, E.,
22 Schwab, E., Zumstein, G., 1993. An OspA serotyping system for *Borrelia*
23 *burgdorferi* based on reactivity with monoclonal antibodies and *ospA* sequence
24 analysis. J Clin Microbiol 31, 340-350.
- 25 Wittwer, C.T., Reed, G.H., Gundry, C.N., Vandersteen, J.G., Pryor, R.J., 2003. High-
26 resolution genotyping by amplicon melting analysis using LCGreen. Clin Chem
27 49, 853-860.
- 28 Wodecka, B., Skotarczak, B., 2005. First isolation of *Borrelia lusitaniae* DNA from
29 *Ixodes ricinus* ticks in Poland. Scand J Infect Dis 37, 27-34.
- 30 Wormser, G.P., Brisson, D., Liveris, D., Hanincova, K., Sandigursky, S.,
31 Nowakowski, J., Nadelman, R.B., Ludin, S., Schwartz, I., 2008. *Borrelia*
32 *burgdorferi* genotype predicts the capacity for hematogenous dissemination
33 during early Lyme disease. J Infect Dis 198, 1358-1364.
- 34 Wormser, G.P., Liveris, D., Nowakowski, J., Nadelman, R.B., Cavaliere, L.F.,
35 McKenna, D., Holmgren, D., Schwartz, I., 1999. Association of specific subtypes
36 of *Borrelia burgdorferi* with hematogenous dissemination in early Lyme disease.
37 J Infect Dis 180, 720-725.
- 38 Wright, S.A., Lane, R.S., Clover, J.R., 1998. Infestation of the southern alligator
39 lizard (Squamata: Anguidae) by *Ixodes pacificus* (Acari: Ixodidae) and its
40 susceptibility to *Borrelia burgdorferi*. J Med Entomol 35, 1044-1049.
- 41 Xu, G., Fang, Q.Q., Keirans, J.E., Durden, L.A., 2003. Molecular phylogenetic
42 analyses indicate that the *Ixodes ricinus* complex is a paraphyletic group. J
43 Parasitol 89, 452-457.
- 44 Younsi, H., Sarih, M., Jouda, F., Godfroid, E., Gern, L., Bouattour, A., Baranton, G.,
45 Postic, D., 2005. Characterization of *Borrelia lusitaniae* isolates collected in
46 Tunisia and Morocco. J Clin Microbiol 43, 1587-1593.

1 **Table 1** List of putative and named species within the LB group spirochetes, their
2 host and vector range and distribution.

Species (c/p) ^a (type strain)	Distribution	Host range	main vector	References for spec. description
<i>B. afzelii</i> (c) (VS461)	Europe, Asia	<i>Apodemus</i> spp, <i>Myodes glareolus</i> , <i>Sorex</i> spp, <i>Sciurus</i> spp, <i>Erinaceus</i> spp, <i>Rattus</i> spp	<i>Ixodes ricinus</i> , <i>I. persulcatus</i> , <i>I. hexagonus</i>	Canica et al. 1993
<i>B. americana</i> (p) (SCW-41)	North America	<i>Thryothorus</i> <i>ludovicianus</i> , <i>Pipilo</i> <i>erythrophthalmus</i>	<i>I. pacificus</i> , <i>I. minor</i>	Rudenko et al. 2009b
<i>B. andersonii</i> (c) (21038)	North America	<i>Sylvilagus</i> spp, (Passeriformes spp)	<i>I. dentatus</i>	Marconi et al 1995
<i>B. bavariensis</i> (p) (PBi)	Europe, Asia (?)	<i>Apodemus</i> spp, <i>Myodes</i> sp, <i>Microtus</i> spp.	<i>I. ricinus</i> , <i>I.</i> <i>persulcatus</i> (?)	Margos et al. 2009
<i>B. bissettii</i> (c) (DN127-cl9-2)	North America, Europe	<i>Neotoma</i> spp, <i>Peromyscus</i> spp, <i>Sigmodon</i> spp EU: unknown	<i>I. pacificus</i> , <i>I. spinipalpis</i> , <i>I. affinis</i> , EU: unknown	Postic et al. 1998
<i>B. burgdorferi</i> (c) (B31)	North America, Europe	<i>Peromyscus</i> spp, <i>Tamias</i> spp, <i>Neotoma</i> spp, <i>Sorex</i> spp, <i>Sciurus</i> spp, <i>Sigmodon</i> spp <i>Erinaceus</i> spp, <i>Rattus</i> spp, <i>Procyon lotor</i> , <i>Turdus</i> <i>migratorius</i> ,	<i>I. ricinus</i> , <i>I. hexagonus</i> , <i>I. scapularis</i> , <i>I. pacificus</i> , <i>I. affinis</i> , <i>I. minor</i> , <i>I. spinipalpis</i> , <i>I. muris</i>	Johnson et al. 1984
<i>B. californiensis</i> (c) (CA446)	Western US	<i>Dipodomys</i> <i>californicus</i>	unknown	Postic et al. 2007
<i>B. carolinensis</i> (c) (SCW-22)	Southeast US	<i>P. gossypinus</i> , <i>N. floridana</i>	unknown (<i>I. minor</i> ?)	Rudenko et al. 2009a
<i>B. garinii</i> (c) (20047)	Europe, Asia, Artic-Antartic circles	<i>Turdus merula</i> , <i>T. philomelos</i> , <i>Parus major</i> , seabirds (Puffin, Guillemot, Kittiwake, Razorbill)	<i>I. ricinus</i> , <i>I. persulcatus</i> , <i>I. uriae</i>	Baranton et al. 1992
<i>B. japonica</i> (c) (HO14)	Japan	<i>Sorex</i> <i>unguiculatus</i> , <i>Apodemus</i> spp, <i>Eothenomys smithi</i>	<i>I. ovatus</i>	Kawabata et al. 1993, Postic et al. 1993
<i>B. kurtenbachii</i> (p) (25015)	Northamerica, (Europe?)	<i>Microtus</i> <i>pennsylvanicus</i> , <i>Zapus hudsonius</i> <i>Peromyscus</i> ?	unknown (<i>I. scapularis</i> ?)	Margos et al. 2010
<i>B. lusitaniae</i> (c) (PoTiB2)	Mediterranean basin	Lacertidae	<i>I. ricinus</i>	LeFleche et al. 1997
<i>B. sinica</i> (c) (CMN3)	China	<i>Niviventer</i> <i>confucianus</i>	<i>I. ovatus</i>	Masuzawa et al. 2001
<i>B. spielmanii</i> (c)	Europe	<i>Glis glis</i> ,	<i>I. ricinus</i>	


(PC-Eq17N5)		<i>Eliomus quercinus</i>		Richter et al. 2006
<i>B. tanukii</i> (c) (Hk501)	Japan	<i>Apodemus</i> sp, <i>Clethrionomys</i> <i>rufocanus</i> , <i>Eothenomys</i> <i>smithii</i>	<i>I. tanuki</i>	Fukunaga et al. 1996
<i>B. turdi</i> (c) (Ya501)	Japan	<i>Turdus</i> spp	<i>I. turdus</i>	Fukunaga et al. 1996
<i>B. valaisiana</i> (c) (VS116)	Europe, Japan	<i>Turdus merula</i> , <i>T. philomelos</i> , <i>Parus major</i>	<i>I. ricinus</i> , <i>I. columnae</i>	Wang et al. 1997
<i>B. yangtze</i> (p) (nd)	China	<i>Niviventer</i> <i>fulvescens</i> , <i>Apodemus</i> sp	<i>I. granulatus</i> , <i>I. nipponensis</i>	Chu et al. 2008
Genomospecies2	United States	unknown	<i>I. spinipalpis</i> , <i>I. pacificus</i>	Postic et al. 2007

^ac – confirmed ; p – proposed; nd = not determined

Table 2 Typing schemes for LB spirochetes using multiple loci

Type of Loci	Loci	purpose	data	reference
chromosomal housekeeping genes	<i>clpA</i> , <i>clpX</i> , <i>nifS</i> , <i>pepX</i> , <i>pyrG</i> , <i>recG</i> , <i>rplB</i> , <i>uvrA</i>	taxonomy, population studies, evolutionary studies	borrelia.mlst.net, >1,200 strains, 327 STs,	Margos et al. 2008, 2009, 2010; Hoen et al. 2009, Ogden et al. 2010, Vollmer et al. 2011, Ogden et al. 2011, Takano et al. 2011
plasmid-encoded Osp, chromosomal: rRNA, intergenic spacer, housekeeping gene	<i>ospA</i> , 16S, <i>p66</i> , 23S-5S IGS, <i>flaB</i>	taxonomy	GenBank ~110 strains	Rudenko et al. 2009, 2010
plasmid-encoded Osp, chromosomal: rRNA, intergenic spacer, housekeeping genes	<i>ospA</i> , 16S, 23S-5S IGS, <i>groEL</i> , <i>hbb</i> , <i>fla</i> , <i>recA</i>	taxonomy	~130 strains	Richter et al. 2006, Postic et al. 2007, Chu et al. 2008
17 plasmid-encoded loci, chromosomal: housekeeping gene	<i>lp54</i> , <i>cp26</i> , <i>cp9</i> , <i>lp17</i> , <i>lp25</i> , <i>lp28-2</i> , <i>lp28-4</i> , <i>lp38</i> , BB0082	population studies	GenBank, ~60 strains	Qiu et al. 2004
plasmid-encoded Osp's, chromosomal: membrane protein, intergenic spacer	<i>ospA</i> , <i>ospC</i> , <i>p66</i> , 16S-23S IGS	population studies	GenBank, ~115 strains	Bunikis et al. 2004, Humphry et al. 2010 (except <i>p66</i>)

1




2

3 **Figure 1.** Factors impacting the evolutionary ecology of LB spirochetes. Biotic
 4 factors are shown next to the host-vector-spirochete triangle. Abiotic factors (such as
 5 climate or landscape) act indirectly on LB spirochetes by impacting on host and
 6 vector populations. The contemporary picture is further compounded by the
 7 evolutionary and demographic history of hosts, vectors, and pathogens.

8

9



1

2 **Figure 2.** Bayesian phylogenetic inferences generated using MLST housekeeping
3 genes (A) and ospC (B) sequences. Previously assigned species are color coded as
4 follows: *B. burgdorferi* s.s. – ■, *B. afzelii* – ◻, *B. garinii* – ●, *B. bavariensis* – ▲,

1 *B. valaisiana* – ○ , and *B. lusitaniae* – ▼ . The MLST tree was rooted with
2 sequences of the relapsing fever spirochetes *B. duttonii*, *B. hermsii*, and *B. turicatae*.
3 The branch length of the outgroup is not according to scale as indicated by slashes.
4 While in the MLST tree LB species cluster monophyletically, this is not the case
5 using *ospC* sequences (original figure A from Population Biology of Lyme
6 Borreliosis Spirochetes; Kurtenbach et al [2010],
7 DOI: 10.1002/9780470600122.ch12; Copyright (2010, John Wiley & Sons); reprinted
8 with permission of John Wiley & Sons, Inc.; original figure B Margos et al. [2009],
9 doi 10.1128/AEM.00116-09, reproduced and modified with permission from the
10 American Society for Microbiology)
11





Figure 3. Multi Locus Sequence Typing. Targetted PCR is used to amplify several genes distributed throughout the genome exhibiting nearly neutral variation. Internal fragments, kept in-frame, of similar length for each gene are used. For each individual gene, fragments of identical length are aligned and compared to sequences in a ‘virtual strain collection centre’, a MLST database, and to each other permitting determination of an allelic profile for each strain. The allelic profile determines the sequence type (ST) and it can be used to infer relationships of descent within bacterial species based on models of clonal expansion and diversification. Concatenated sequences of all genes can be used for phylogenetic inferences. The accumulative nature of MLST database makes it an attractive instrument to understand intra- and inter-specific relationships of bacteria.

1
2



3

4 **Figure 4.** Map showing the global distribution of the LB species. The shaded areas
5 show the distribution of tick vectors. Seven species of LB group spirochetes are found
6 in North America, eight species in Europe, and eight species in Asia, two species
7 overlap in the Old and New Worlds, three in Europe and Asia (see text for details).




Figure 5. goeBURST diagrams based on the multi-locus allelic profiles for *B. garinii* (A), *B. valaisiana* (B) and *B. afzelii* (C). Each coloured box represents an ST. The colour and size of the boxes corresponds to geographic region and the number of that ST found. STs unique to a particular country were coloured as follows: red England, blue France, yellow Germany, green Latvia, purple Scotland. Those STs that were found in more than one country are grey. STs connected by black or blue lines are single-locus variants (SLVs) and STs connected by grey or green lines are double-locus variants (DLVs) (original figure from Vollmer et al. [2011] *Environmental Microbiology*, doi:10.1111/j.1462-2920.2010.02319.x, reproduced with permission from John Wiley and Sons)





Figure 6 Proposed post-glacial migration routes for three small mammal species taken from Hewitt (1999) based on fossil and molecular data. (original figure from Hewitt [1999] *Biological Journal of the Linnean Society*, doi:10.1111/j.1095-8312.1999.tb01160.x, partially reproduced with permission from John Wiley and Sons).

1



2

3 **Figure 7.** A population snapshot of 244 samples of *Borrelia burgdorferi* found in
 4 Canada (166 samples) and the Unites States (78 samples) as determined by spatial
 5 analysis using spatialepidemiology.net. The figure reveals correspondence of
 6 sequence type and geographic distribution. Most ST were found either in the
 7 Northeast or the Midwest suggesting limited gene flow between populations.

8

9




Figure 8 Neighbour joining tree generated using concatenated sequences of MLSA housekeeping genes showing LB groups species. Black dots indicate species that occur in North America, circles indicate species that occur in Eurasia, grey dots indicate species that occur in the Old and New Worlds. The scale bar shows 1 % divergence. Branch confidence values calculated using a bootstrap procedure with 100 repetitions (original figure from Margos et al. [2010] *Ticks and Tick-borne Diseases*, doi: 10.1016/j.ttbdis.2010.09.002, modified and reproduced with permission from Elsevier)




Figure 9 Graphic representation of the ‘time’ captured by various genetic elements used for typing of bacterial microorganisms. The highly conserved 16S locus reveals deep evolutionary relationships but is unable to capture recent events. Fast evolving genetic elements, such as loci under diversifying selection, microsatellites or variable number of tandem repeats (VNTR) may reveal very recent events but – due to saturation – are not able to ‘see’ ancient events. Intergenic spacer (IGS) regions are supposed to be selectively neutral and should therefore accumulate mutations indiscriminately and linear to time. IGS may be short and saturate quickly or may contain regulatory elements which might not permit all mutations to be fixed. Due to the slow evolution of housekeeping genes multilocus sequence typing captures the intermediate relationship of bacteria. Genome-wide SNPs provide the broadest ‘view’ on an organism past as these are able to capture recent as well as ancient events.